

**This lecture:**

- More on convex optimization
- Minima of convex problems
- First and second order characterizations of convex functions
- Least squares revisited
- Strict convexity and uniqueness of optimal solutions
- Optimality condition for convex problems

Instructor:  
Amir Ali Ahmadi  
Fall 2014

TAs: Y. Chen,  
G. Hall,  
J. Ye

Recall from the last lecture that a convex optimization problem is a problem of the form:

$$\begin{array}{ll} \min. & f(x) \\ \text{s.t.} & g_i(x) \leq 0 \quad i = 1, \dots, m \\ & h_i(x) = 0 \quad i = 1, \dots, k \end{array}$$

where

- Each  $h_i$  is affine:  $h_i(x) = a_i^T x - b_i$
- $f, g_1, \dots, g_m$  are convex:  $f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y)$

$$\forall x, y \in \text{dom}(f), \forall \lambda \in [0, 1]$$

Similarly for the  $g_i$ 's.

- Today we start off by proving results that explain why we give special attention to convex optimization problems.
  - In a convex problem, every local minimum is automatically a global minimum. (This is true even for the more abstract definition of a convex optimization problem from the last lecture that only required the feasible set to be a convex set.)
  - In the unconstrained case, every stationary point (i.e., zero of the gradient) is automatically a global minimum.
- We will also see new characterizations for convex functions that make the task of checking convexity somewhat easier, though in general checking convexity can be a very difficult problem [AOPT13].

Let's recall the definition of local and global minima and generalize them to the constrained setting.

$$\begin{array}{ll} \min. & f(x) \\ \text{s.t.} & x \in \Omega \end{array} \quad (\text{e.g., } \Omega = \{x \mid g_i(x) \leq 0, h_j(x) = 0\})$$

**Definition:** A point  $x^* \in \mathbb{R}^n$  is

- *feasible*, if  $x^* \in \Omega$ ; i.e.,  $g_i(x^*) \leq 0, \forall i, h_j(x^*) = 0, \forall j$
- a *local minimum*, if feasible, and if  $\exists \delta > 0$  s.t.


$$f(x^*) \leq f(x), \forall x \text{ s.t. } x \in \Omega \text{ and } \|x - x^*\| < \delta$$

- a *strict local minimum*, if feasible, and if  $\exists \delta > 0$  s.t.

$$f(x^*) < f(x), \forall x \neq x^* \text{ s.t. } x \in \Omega \text{ and } \|x - x^*\| < \delta$$

- a *global minimum*, if feasible, and if  $f(x^*) \leq f(x), \forall x \in \Omega$
- a *strict global minimum*, if feasible, and if  $f(x^*) < f(x), \forall x \in \Omega, x \neq x^*$

- $x_1$ : not feasible
- $x_2$ : strict global minimum
- $x_3$ : feasible
- $x_4$ : strict local minimum



Our next few theorems show the nice features of convex problems in terms of inferring global properties from local ones.

**Theorem.** Consider an optimization problem

$$\begin{array}{ll} \min. & f(x) \\ \text{s.t.} & x \in \Omega \end{array}$$

where  $f$  is a convex function and  $\Omega$  is a convex set. Then, every local minimum is also a global minimum.

**Proof.**

Let  $x$  be a local minimum. Suppose for the sake of contradiction that  $x$  is not a global minimum.

$$\Rightarrow \exists y \in \Omega, \text{ s.t. } f(y) < f(x).$$

But  $x \in \Omega, y \in \Omega, \Omega$  convex  $\Rightarrow \lambda x + (1 - \lambda)y \in \Omega, \forall \lambda \in [0,1]$

and  $f$  convex  $\Rightarrow f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y)$

$$< \lambda f(x) + (1 - \lambda)f(x) = f(x), \forall \lambda \in [0,1].$$

As  $\lambda \rightarrow 1$ ,  $(\lambda x + (1 - \lambda)y) \rightarrow x$ . So there are points arbitrarily close to  $x$  with a better objective value than  $x$ . This contradicts local optimality of  $x$ .  $\square$

**Intuition:**



$$f(y) < f(x) \Rightarrow f(\text{all line}) < f(x)$$


## First and second order characterization of convex functions

**Theorem.** Suppose  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  is twice differentiable over its domain. Then, the following are equivalent:

- (i)  $f$  is convex.
- (ii)  $f(y) \geq f(x) + \nabla f^T(x)(y - x), \quad \forall x, y \in \text{dom}(f).$
- (iii)  $\nabla^2 f(x) \geq 0, \quad \forall x \in \text{dom}(f).$

### Interpretation:


- (i) The first order Taylor expansion at any point is a global under estimator of the function



- (ii) Function has *nonnegative curvature* everywhere:

"It curves up".

- In one dimension:  
 $f''(x) \geq 0, \forall x \in \text{dom}(f)$



We prove  $(i) \Leftrightarrow (ii)$ . For  $(ii) \Leftrightarrow (iii)$ , see, e.g., Theorem 22.5 of [CZ13].

**Proof:** ([Tit13])(i)  $\Rightarrow$  (ii) If  $f$  convex, by definition

$$f(\lambda y + (1-\lambda)x) \leq \lambda f(y) + (1-\lambda)f(x), \quad \forall \lambda \in [0,1], x, y \in \text{dom}(f).$$

After rewriting, we have

$$\begin{aligned} f(x + \lambda(y-x)) &\leq f(x) + \lambda(f(y) - f(x)) \\ \Rightarrow f(y) - f(x) &\geq \frac{f(x + \lambda(y-x)) - f(x)}{\lambda}, \quad \forall \lambda \in (0,1]. \end{aligned}$$

As  $\lambda \downarrow 0$ , we get

$$f(y) - f(x) \geq \nabla f^T(x) (y-x). \quad (1)$$

(ii)  $\Rightarrow$  (i) Suppose (1) holds  $\forall x, y \in \text{dom}(f)$ .Take any  $x, y \in \text{dom}(f)$  and let  $z = \lambda x + (1-\lambda)y$ .

We have

$$f(x) \geq f(z) + \nabla f^T(z) (x-z) \quad (2)$$

$$f(y) \geq f(z) + \nabla f^T(z) (y-z). \quad (3)$$

Multiplying (2) by  $\lambda$ , (3) by  $(1-\lambda)$  and adding, we get

$$\begin{aligned} \lambda f(x) + (1-\lambda)f(y) &\geq f(z) + \nabla f^T(z) (\lambda x + (1-\lambda)y - z) \\ &= f(z) \\ &= f(\lambda x + (1-\lambda)y). \quad \square \end{aligned}$$

**Corollary.** Consider an unconstrained optimization problem:

$$\begin{aligned} \min. & f(x) \\ \text{s.t. } & x \in \mathbb{R}^n, \end{aligned}$$

where  $f$  is convex and differentiable. Then, any point  $\bar{x}$  that satisfies  $\nabla f(\bar{x}) = 0$ , is a global minimum.

**Proof.** From the first order characterization of convexity

we have 
$$f(y) \geq f(x) + \nabla f^T(x)(y - x) \quad \forall x, y$$

In particular, 
$$f(y) \geq f(\bar{x}) + \nabla f^T(\bar{x})(y - \bar{x}) \quad \forall y$$

Since  $\nabla f(\bar{x}) = 0$ , we get

$$f(y) \geq f(\bar{x}) \quad \forall y. \quad \square$$

**Remark 1.** Recall that  $\nabla f(x) = 0$  is always a necessary condition for local optimality in an unconstrained problem. The theorem says that for convex problems  $\nabla f(x) = 0$  is not only necessary, but also sufficient for local *and global* optimality.

**Remark 2.** Recall that in absence of convexity,  $\nabla f(x) = 0$  is not sufficient even for local optimality (e.g., think of  $f(x) = x^3$  and  $\bar{x} = 0$ ).

**Remark 3.** Recall that another necessary condition for (unconstrained) local optimality of a point  $x$  was:  $\nabla^2 f(x) \succcurlyeq 0$ .

- Note that a convex function automatically passes this test.

## Quadratic functions revisited

- Let  $f(x) = x^T A x + b x + c$  ( $A$  symmetric)
- When is  $f$  convex?
  - Let's use the second order test:

$$\nabla^2 f(x) = 2A$$

$$\Rightarrow f \text{ convex} \Leftrightarrow A \succeq 0 \text{ (independent of } b, c)$$

- Consider the unconstrained optimization problem

$$\min_x x^T A x + b x + c$$

- $A \not\succeq 0$  ( $f$  not convex)  $\Rightarrow$  unbounded below (why?)
- $A \succ 0 \Rightarrow$  convex (in fact,  $\Rightarrow f$  strictly convex as we see next)

$$\text{Unique solution: } x^* = -\frac{1}{2} A^{-1} b \text{ (why?)}$$


- $A \succeq 0 \Rightarrow f$  convex. Optimal value bounded, but there could be many optimal solutions.

Let  $\bar{x}$  be an eigen vector, with a negative eigen value  $\lambda$ .


$$A \bar{x} = \lambda \bar{x} \Rightarrow \bar{x}^T A \bar{x} = \lambda \bar{x}^T \bar{x} < 0.$$

$$f(\alpha \bar{x}) = \alpha^2 \underbrace{\bar{x}^T A \bar{x}}_{< 0} + \alpha b \bar{x} + c$$


$$f(\alpha \bar{x}) \rightarrow -\infty \text{ as } \alpha \rightarrow \infty.$$



$$A \succ 0$$



$$A \succeq 0 \\ \text{(but } A \not\succ 0)$$



$$A \not\succeq 0$$





## Least squares, revisited.

Given:  $A$   $m \times n$  matrix (Assume columns of  $A$  are linearly independent)  
 $b$   $m \times 1$  vector

Solve:  $\min_x \|Ax - b\|^2$

Let  $f(x) = \|Ax - b\|^2 = (Ax - b)^T (Ax - b)$   
 $= x^T A^T A x - 2x^T A^T b + b^T b.$

Strictly

Convex!


(b/c  $A^T A \succ 0$ )

$$\nabla f(x) = 2A^T A x - 2A^T b$$

Called

$$\nabla f(x) = 0 \Rightarrow A^T A x = A^T b \leftarrow \text{"Normal Equations"}$$

$$\Rightarrow x = (A^T A)^{-1} A^T b$$



$A^T A$  is invertible b/c its null space is just the origin:

$$A^T A x = 0 \Rightarrow x^T A^T A x = 0 \Rightarrow (Ax)^T (Ax) = 0 \Rightarrow \|Ax\|^2 = 0 \Rightarrow Ax = 0$$

$$\Rightarrow x = 0.$$

Columns of  $A$  linearly independent.

$$\nabla^2 f(x) = 2A^T A \succ 0 \quad (\text{b/c } x^T A^T A x = \|Ax\|^2 \geq 0 \text{ and } = 0 \Leftrightarrow x = 0)$$

$\Rightarrow x = (A^T A)^{-1} A^T b$  is a strict local minimum.  $\square$

This is the best conclusion we could make before without knowing that  $f$  is convex. Now that we know  $f$  is (strictly) convex, we immediately know that the solution  $x = (A^T A)^{-1} A^T b$  is a (strict) **global** minimum.

## Characterization of Strict Convexity

Recall that we say  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  is strictly convex, if  $\forall x, y, x \neq y, \forall \lambda \in (0,1)$ ,

$$f(\lambda x + (1 - \lambda)y) < \lambda f(x) + (1 - \lambda)f(y).$$

- $f$  strictly convex  $\Rightarrow f$  convex (obvious from the definition)
- $f$  convex  $\not\Rightarrow f$  strictly convex

e.g.,  $f(x) = x^4 \quad (x \in \mathbb{R})$

- Second order sufficient condition:


$$\nabla^2 f(x) > 0 \quad \forall x \in \Omega \Rightarrow f \text{ strictly convex on } \Omega$$

- Converse not true:

$$f(x) = x^4 \quad (x \in \mathbb{R})$$

$f$  is strictly convex (why?).

But  $f''(0) = 0$  (check)



- First order characterization:

$$f \text{ strictly convex on } \Omega \subseteq \mathbb{R}^n$$

$\Leftrightarrow$

$$f(y) > f(x) + \nabla f^T(x)(y - x), \quad \forall x, y \in \Omega, x \neq y$$

- One of the main uses of strict convexity is to ensure uniqueness of optimal solutions. We see this next.

## Strict Convexity and Uniqueness of Optimal Solutions

**Theorem.** Consider an optimization problem

$$\begin{array}{ll} \min. & f(x) \\ \text{s.t.} & x \in \Omega \end{array}$$

where  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  is *strictly convex* on  $\Omega$  and  $\Omega$  is a convex set. Then, the (optimal) solution is unique (assuming it exists).

**Proof.** Suppose there were two optimal solutions  $x, y \in \mathbb{R}^n$ . This means that  $x, y \in \Omega$  and

$$f(x) = f(y) \leq f(z), \forall z \in \Omega \quad (1)$$

But consider  $z = \frac{x+y}{2}$ . By convexity of  $\Omega$ , we have  $z \in \Omega$ . By strict convexity,

$$\text{we have } f(z) = f\left(\frac{x+y}{2}\right)$$

$$< \frac{1}{2}f(x) + \frac{1}{2}f(y)$$

$$= \frac{1}{2}f(x) + \frac{1}{2}f(x) = f(x).$$

But this contradicts (1).  $\square$

**Practice:** for each function below, determine whether it is convex, strictly convex, or neither.

- $f(x) = (x_1 - 3x_2)^2$
- $f(x) = (x_1 - 3x_2)^2 + (x_1 - 2x_2)^2$
- $f(x) = (x_1 - 3x_2)^2 + (x_1 - 2x_2)^2 + x_1^3$
- $f(x) = |x| \quad (x \in \mathbb{R})$
- $f(x) = ||x|| \quad (x \in \mathbb{R}^n)$

## An Optimality Condition for Convex Problems

**Theorem.** Consider an optimization problem


$$\begin{aligned} \min f(x) \\ \text{s.t. } x \in \Omega, \end{aligned}$$

where  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  is convex and differentiable and  $\Omega$  is a convex. Then, a point  $x$  is optimal if and only if  $x \in \Omega$  and

$$\nabla f(x)^T (y - x) \geq 0, \forall y \in \Omega.$$

- **What does this mean?**

- If you move from  $x$  towards any feasible  $y$ , you will increase  $f$  locally.
- $-\nabla f(x)$  (assuming it is nonzero) serves as a hyperplane that "supports" the feasible set  $\Omega$  at  $x$  (see figure below).



- The necessity of the condition holds independent of convexity of  $f$ .
- Convexity is used in establishing sufficiency.
- If  $\Omega = \mathbb{R}^n$ , can you see how the condition above reduces to our first order unconstrained optimality condition  $\nabla f(x) = 0$ ?
  - Hint: take  $y = x - \nabla f(x)$ .

**Proof.**

(Sufficiency)

Suppose  $x \in \Omega$ satisfies  $\nabla f^T(x)(y-x) \geq 0, \forall y \in \Omega. \quad (1)$ 

By the first order characterization of convexity we have:

$$f(y) \geq f(x) + \nabla f^T(x)(y-x), \quad \forall y \in \Omega \quad (2)$$

$$(1) + (2) \Rightarrow f(y) \geq f(x), \quad \forall y \in \Omega.$$

 $\Rightarrow x$  is optimal.


(necessity)

Suppose  $x$  is optimal, but for some  $y \in \Omega$  we had

$$\nabla f^T(x)(y-x) < 0.$$

Consider  $g(\alpha) := f(x + \alpha(y-x))$ .Because  $\Omega$  is convex,  $\forall \alpha, x + \alpha(y-x) \in \Omega$ .Observe that  $g'(\alpha) = (y-x)^T \nabla f(x + \alpha(y-x))$ .

$$\Rightarrow g'(0) = (y-x)^T \nabla f(x) < 0.$$



$$\Rightarrow \exists \delta > 0 \text{ s.t. } g(\alpha) < g(0) \quad \forall \alpha \in (0, \delta).$$

$$\Rightarrow f(x + \alpha(y-x)) < f(x) \quad \forall \alpha \in (0, \delta).$$

But this contradicts

optimality of  $x$ .  $\square$

### Notes:

- Further reading for this lecture can include the first few pages of Chapters 2,3,4 of [BV04]. Your [CZ13] book defines convex sets in Section 4.3. Convex optimization appears in Chapter 22. The relevant sections are 22.1-22.3.

### References:

- [AOPT13] A.A. Ahmadi, A. Olshevsky, P.A. Parrilo, and J.N. Tsitsiklis. NP-hardness of checking convexity of quartic forms and related problems. *Mathematical Programming*, 2013.  
[http://web.mit.edu/~a\\_a/Public/Publications/convexity\\_nphard.pdf](http://web.mit.edu/~a_a/Public/Publications/convexity_nphard.pdf)
- [BV04] S. Boyd and L. Vandenberghe. *Convex Optimization*. Cambridge University Press, 2004.  
<http://stanford.edu/~boyd/cvxbook/>
- [CZ13] E.K.P. Chong and S.H. Zak. *An Introduction to Optimization*. Fourth edition. Wiley, 2013.
- [Tit13] A.L. Tits. *Lecture notes on optimal control*. University of Maryland, 2013.