
CAUSES AND CONSEQUENCES OF EARLY-LIFE HEALTH*

ANNE CASE AND CHRISTINA PAXSON

We examine the consequences of child health for economic and health outcomes in adulthood, using height as a marker of childhood health. After reviewing previous evidence, we present a conceptual framework that highlights data limitations and methodological problems that complicate the study of this topic. We then present estimates of the associations between height and a range of outcomes—including schooling, employment, earnings, health, and cognitive ability—measured in five data sets from early to late adulthood. These results indicate that, on average, taller individuals attain higher levels of education. Height is also positively associated with better economic, health, and cognitive outcomes. These associations are only partially explained by the higher average educational attainment of taller individuals. We then use data from the National Longitudinal Survey of Youth 1979 Children and Young Adults survey to document the associations between health, cognitive development, and growth in childhood. Even among children with the same mother, taller siblings score better on cognitive tests and progress through school more quickly. Part of the differences found between siblings arises from differences in their birth weights and lengths attributable to mother's behaviors while pregnant. Taken together, these results support the hypothesis that childhood health influences health and economic status throughout adulthood.

Numerous studies have examined the association between economic status (as measured by income, occupation, or education) and health. Despite occasional anomalies, three general patterns emerge from the data. The first is that health is strongly related to economic status: wealthier people live longer and have lower morbidity. This is true whether one looks across countries, within countries at any point in time, or over time with economic growth. The second is that the association between health and wealth—the “gradient”—is evident throughout the income distribution. The gradient does not merely reflect the association between poor health and absolute poverty; it is evident in wealthy countries and among wealthier individuals within countries. The third is that the association between health and wealth is found across a wide range of ages. Differences in health across economic groups are apparent at birth: newborns from less-well-off families or with less-well-educated mothers have higher rates of mortality and lower birth weights than more-advantaged newborns. The association between health and wealth becomes more pronounced until early old age but then eventually declines, possibly the result of selective mortality.

These patterns, although uncontroversial, have been subject to many interpretations. One hypothesis is that poor health is the product of low socioeconomic status. This may be due to the effects of material deprivation—less-nutritious food, lower-quality medical care, and lower-quality housing—on health. Relatedly, those with lower income or less education may live in more stressful environments, which could lead to greater physiological “wear and tear” on metabolic and immune systems and to a greater reliance on unhealthy behaviors, such as smoking (Adler et al. 1994.) An alternative view is that poor health reduces

*Anne Case and Christina Paxson, Center for Health and Wellbeing, Princeton University. Direct correspondence to Anne Case, 367 Wallace Hall, Princeton University, Princeton, NJ 08544; e-mail: accase@Princeton.edu. This article was written for the RAND 15th Anniversary Gala held on July 8–9, 2009. We thank conference participants and two anonymous referees for useful comments, and Kimberly Bryan for expert research assistance. We gratefully acknowledge funding from the National Institutes of Health under the Demography of Aging Center Grant P30 AG024361.

economic status. Children in poor health may have lower educational attainment, which reduces their adult earnings potential. Adults with health problems may work less or may be less productive on their jobs, leading to lower earnings, and they may be subject to greater out-of-pocket medical costs, reducing asset accumulation.

These two views need not be mutually exclusive. Indeed, it is likely that there is a lifelong interplay between health and wealth, with different factors coming into prominence at different stages of the life cycle. Furthermore, the relationships between health and socioeconomic status are certainly dynamic—health at one stage of life has implications for future socioeconomic status, and socioeconomic status influences future health. In this article, we focus on a specific aspect of the health-wealth gradient: the association between childhood health, on one hand, and economic and health outcomes later in life, on the other.

Educational attainment is an especially important adult outcome to consider. Schooling is obtained throughout childhood and has a large influence on earnings, which may in turn affect health. Education may also have a direct effect on health and health behaviors later in life (Cutler and Lleras-Muney 2008). Researchers have documented many dimensions of the association between childhood health and educational attainment.¹ In many large, cross-sectional studies, researchers have shown that cohorts affected by health insults *in utero* have lower educational attainment, on average, than do preceding and following birth cohorts that did not face the same health insult. Using a variety of data sets, researchers have also documented the association between low birth weight and educational attainment. In addition, several birth cohort studies allow researchers to trace the impact of childhood health (measured using birth weight, mothers' assessments of children's health, chronic conditions in childhood, days spent in bed, and hospitalization episodes) on progress through school and educational attainment.

Estimates of the importance of health *in utero* and during childhood on educational attainment vary but are often found to be large and significant. Almond (2006), for example, estimated that children born to women infected with influenza during the 1918 pandemic received, on average, five fewer months of education than children born before or conceived after this event. Case, Fertig, and Paxson (2005) found, using data from the 1958 British birth cohort study, that children born at low birth weight passed 0.5 fewer O-level exams (qualifying exams taken in British secondary schools prior to 1988) than other children, on average. This effect is larger than that predicted for moving a cohort member's household from the 90th to the 10th percentile of family income at age 16. Smith (2009), using retrospective reports on childhood health in the PSID, found a large and significant association between health reports and educational attainment in OLS regressions but not in regressions that included maternal fixed effects. In contrast, Magnusson, Rasmussen, and Gyllenstein (2006) found that the effect of height on the probability of obtaining higher education among men in Sweden is almost as large within sibships (odds ratio [OR] of 1.08 per 5 cm increase in height) as between any two Swedish men (OR = 1.10). Black, Devereux, and Salvanes (2007), using data on Norwegian twins, found that within twin pairs birth weight is a significant predictor of both IQ at age 18 and high school completion, with fixed-effect estimates similar in size to cross-sectional estimates.

Less has been written about the impact of early life health on transitions to the labor force, employment, and earnings in young adulthood. This is in part because studies following individuals from birth through middle age are still relatively scarce. Much of the evidence that we have comes again from being able to identify the impact of a large health shock on a birth cohort (e.g., the 1918 influenza pandemic) or from a handful of panel studies, including the British birth cohorts of 1958 and 1970, called the National Child

1. See Currie (2009) for a thorough review of the relationships between health in childhood and a range of outcomes, including education.

Development Study (NCDS) and the British Cohort Study (BCS), respectively. Case et al. (2005) found that for men followed in the NCDS, markers of early-life health make a large incremental contribution to the pseudo- R^2 of being employed at age 42. Specifically, with controls for family background, including the ages at which mother and father left school and father's and grandfathers' social classes, they found that chronic conditions in childhood and height at age 16 contribute 0.25 to the pseudo- R^2 in a probit regression of employment at age 42. Black et al. (2007) found, within twin pairs in Norway, that the twin born at higher birth weight reported significantly higher earnings in early adulthood.

To date, life course models have had to rely on data collected from birth to middle age (as in the 1958 and 1970 British cohort studies) or from middle age to old age and death (as in the U.S. Health and Retirement Study and in the U.K. Whitehall II Study). Longitudinal studies that follow cohort members from cradle to grave are, at this point, rare. An exception is the Helsinki Birth Cohort Study, which David Barker, Johann Eriksson, and colleagues have used to examine the fetal origins hypothesis. Beyond this, estimates of the extent to which early-life health affects socioeconomic status and health in old age are based primarily on retrospective questions asked in surveys currently being conducted on older adults.²

With a lack of information on health in childhood, researchers have begun to use height as a marker of early life health and nutrition, which serves as a bridge from health in childhood to health and socioeconomic status at older ages.³ Our article follows in this relatively new line of research. In the next section, we present a conceptual framework that captures the dynamic associations between health and economic status over the life course. The conceptual framework is used to highlight the methodological issues and data demands that complicate research on the long-run effects of childhood health and to motivate the use of height as a bridge from childhood into adulthood. We then present evidence on the associations between height and a range of outcomes in adulthood, including educational attainment, employment, earnings, health, and cognitive function. Finally, we consider the question of what aspects of the early-life environment height captures. Using data from the National Longitudinal Survey of Youth 1979 (NLSY79) Children and Young Adults study, we show that height is related to a variety of children's health and developmental outcomes, including health at birth and cognitive development within childhood.

CONCEPTUAL FRAMEWORK

The framework we discuss here, which follows that presented in Case et al. (2005), elucidates mechanisms through which early-life experiences may influence later-life outcomes. For simplicity, we assume there are three stages in life: childhood, young adulthood, and older adulthood. We also collapse *economic status* and *health status* into single variables, although each could be thought of as containing several components. For example, *economic status* could include earnings, asset income, and transfer income; *health* could include chronic conditions, acute conditions, and cognitive function. Distinguishing between these different components may be important for empirical work, but not for laying out the framework.

We express economic status in older adulthood (E_A) and a measure of health in older adulthood (H_A) as linear functions of indicators of economic status and health in young adulthood (E_Y and H_Y respectively), educational attainment (S), and measures of economic

2. Smith (2009), for example, found both in ordinary least squares and maternal fixed-effect estimation that retrospective reports of health in childhood have significant effects on earnings and wealth in adulthood in the PSID.

3. For many studies, height is the only marker available for childhood health. We recognize that there are many aspects of childhood health that do not work through height. In the NCDS, Case et al. (2005) found, independent of height, that the number of chronic conditions a cohort member faced at ages 7 and 11 have large and significant associations with educational attainment.

status and health in childhood (E_C and H_C). Health in childhood can include prenatal health, nutrition and illnesses in childhood, and cognitive development.

$$E_A = \beta_0 + H_Y\beta_{HY} + E_Y\beta_{EY} + S\beta_S + H_C\beta_{HC} + E_C\beta_{EC} + \varepsilon_A^E. \quad (1)$$

$$H_A = \alpha_0 + H_Y\alpha_{HY} + E_Y\alpha_{EY} + S\alpha_S + H_C\alpha_{HC} + E_C\alpha_{EC} + \varepsilon_A^H. \quad (2)$$

Economic status, health, and educational attainment in young adulthood are assumed to be functions of economic and health status in childhood:

$$E_Y = \gamma_0 + H_C\gamma_{HC} + E_C\gamma_{EC} + \varepsilon_Y^E. \quad (3)$$

$$H_Y = \delta_0 + H_C\delta_{HC} + E_C\delta_{EC} + \varepsilon_Y^H. \quad (4)$$

$$S = \theta_0 + H_C\theta_{HC} + E_C\theta_{EC} + \varepsilon^S. \quad (5)$$

Eqs. (1) to (5), taken together, are general enough to be consistent with many hypotheses concerning the relationships between health and economic status at different stages of the life cycle. Much of the literature on health and economic status has been concerned with estimating Eqs. (1) and (2), with a focus on the “cross effects” of adult health on later economic status and of adult economic status on later-life health. One hypothesis (most closely associated with the work of Michael Marmot and colleagues) using data on British civil servants, is that the economic gradient in health observed among adults is largely due to the effects of economic status (defined in the Whitehall II study by civil service grade) on disease risk ($\alpha_{EY} \neq 0$). (See, e.g., Brunner et al. 1999; Chandola et al. 2003; and Marmot et al. 2001.) An alternative hypothesis is that the gradient in adulthood reflects the effects of poor health on earnings and assets ($\beta_{HY} \neq 0$).

Our focus here is on the effects on childhood health on later-life outcomes. Eqs. (3), (4), and (5) indicate that childhood health and economic status determine the human capital that individuals carry into young adulthood in the form of higher earnings potential, better health, and greater levels of schooling. (In Eq. (3), the effect of schooling on economic status has been subsumed in the effects of health and economic status in childhood.) The effect of childhood circumstances on later adulthood depends on the parameters in (1) and (2). Childhood circumstances may have both direct effects—even controlling for educational attainment, economic status, and health in young adulthood—and indirect effects that operate through the outcomes in young adulthood. Substitution of Eqs. (3), (4), and (5) into Eqs. (1) and (2) yields reduced-form equations for health and economic status in later adulthood, expressed solely as functions of childhood characteristics. Estimates of these reduced-form equations provide information on the combined direct and indirect effects of childhood circumstances.

The data demands for estimating Eqs. (1) through (5) are high: the model requires data that follow individuals from early childhood through later life. Few such data sets exist. However, if there are no direct effects of childhood health and economic status on outcomes in older adulthood (i.e., $\beta_{HC} = \beta_{EC} = \alpha_{HC} = \alpha_{EC} = 0$), then Eqs. (1) and (2) can be estimated using data sets that follow individuals from younger to older adulthood; provided that accurate measures of education, health, and economic status in young adulthood are available, further information on childhood is unnecessary. Likewise, Eqs. (3), (4), and (5) can be estimated using data that follow individuals from infancy to young adulthood. This “separability” of Eqs. (1) and (2) from Eqs. (3), (4), and (5), if appropriate, greatly eases data demands.

A large body of research relies on the assumption of separability. For example, a discussion of results from the Whitehall II data argued that among British civil servants, “early life circumstances are important because they influence adult social circumstances, which in turn influence disease risk” (Marmot et al. 2001:305). In other work, Brunner et al. (1999:762) stated that “whatever the salient features of the adult socioeconomic environment

may be, it seems they are equally or more important than circumstances in childhood” in determining cardiovascular risk among British civil servants. Although it is convenient to assume that childhood circumstances operate only through their effects on outcomes in early adulthood—especially when working with data that do not contain information on childhood—another body of research argues that this is incorrect. For example, evidence on the well-known fetal origins hypothesis supports the idea that fetal health has a direct effect on health in later life ($\alpha_{HC} \neq 0$). (See, e.g., Barker 1992, 1995.)

Even with data sets that contain information over the lifespan, several econometric issues can complicate estimation of the effects of childhood health on adult outcomes. One issue is that because of the well-documented associations between child health (H_C) and childhood economic circumstances (E_C), it may be difficult to separately identify the effects of these variables on health and economic status in adulthood. If good measures of economic status are unavailable, their effects are likely to load onto measures of child health, and vice versa. This issue will be important when interpreting estimates of the associations between height (as a marker for child health) and outcomes in adulthood. As we discuss below, height is influenced by a wide range of environmental factors experienced in childhood. Although a finding that height is associated with adult outcomes provides support for the hypothesis that childhood circumstances have long-lasting effects, it does not provide information on which aspects of the environment in childhood are important.

A closely related problem is unobserved individual heterogeneity. For example, individuals may have unobserved personality traits or genetic characteristics that result in both poor health and low socioeconomic status throughout the life span. If so, it may appear as if childhood health (possibly measured as height) has effects on adult outcomes, when in fact the association is driven by these unobserved characteristics. There may also be unobservable differences across children in the quality of parenting they receive that is correlated with children’s health and that affects outcomes in adulthood.

To the extent that these traits or genetic characteristics are family-specific, one strategy is to use information on siblings or (in some cases) on twin pairs to control for unobserved heterogeneity. Within-sibling or within-twin estimation techniques have been useful in identifying the impact of early-life circumstance, such as low birth weight, on educational and health outcomes in adulthood (Behrman and Rosenzweig 2004; Black et al. 2007). Within-twin studies have an advantage over sibling studies in that shared aspects of the uterine environment and (with monozygotic twins) genetic endowments can be controlled for. However, there are also disadvantages to twin studies: inferences about singleton births that are drawn from the experiences of twins might not be valid. Furthermore, most nationally representative surveys contain too few twin pairs to use for analysis. In the last section of this article, we rely on within-sibling methods to examine how children’s heights are related to health and development in childhood.

HEIGHT, ECONOMIC STATUS, AND HEALTH STATUS IN ADULTHOOD

This section presents estimates of the associations between height and several measures of economic and health status, estimated using data collected at different points in adulthood in five studies conducted in the United States and the United Kingdom. These results provide information on the role of environmental factors, which influence growth in childhood, in determining health and well-being much later in life. We begin with an overview of the determinants of growth and height, with an emphasis on how environmental conditions *in utero* and during early childhood influence height.

Determinants of Growth and Height

Adult height reflects cumulative growth to maturity. After a period of rapid growth from ages 0 to 3, growth becomes relatively stable at approximately 6 cm per year until adolescence, at which point an adolescent growth spurt accelerates growth to an (adolescent) peak

height velocity of approximately 10 cm per year. Final adult height depends on the timing and duration of the adolescent growth spurt. (See Beard and Blaser [2002] for discussion and references.)

Age-specific growth patterns and final adult height depend on a combination of factors, including genes, environmental conditions (particularly undernutrition and illness), and gene-environment interactions. Although genes are key determinants of individual height, many studies have suggested that differences in average height across populations are due largely to environmental factors (Steckel 1995). The increase in heights throughout the developed world during the twentieth century was too rapid to be due to selection and genetic variation (Beard and Blaser 2002).

The environmental factors thought to influence adult height include intrauterine conditions and conditions in early childhood. Both low birth weight and maternal smoking during pregnancy are significant predictors of lower stature in adulthood (Li, Manor, and Power 2004; Saenger et al. 2007). Adult height is also sensitive to environmental conditions experienced in childhood. Health and nutrition in the period from birth to age 3 are important determinants of adult height. The speed of growth is more rapid during this period than at any other during the life course, and nutritional needs are greatest at this point. Infections (especially gastrointestinal and respiratory) can be frequent and severe in early childhood, and such infections and the body's inflammatory responses to them can impair growth (Crimmins and Finch 2006). In addition, children are at special risk from poor caregiving at youngest ages; once children are more autonomous, they may be better equipped to protect themselves. Research on institutionalized children who are adopted internationally shows that children experience rapid catch-up growth after adoption, although the extent of catch-up declines with the age of adoption (Rutter et al. 1998). These results provide striking evidence of the importance of children's nutritional and psychosocial environments on growth.

Environmental factors that influence height may also affect health throughout life. Undernutrition, chronic infections, and psychosocial stress acting at critical periods of development can induce permanent changes in body organs and systems. These changes, initiated and "programmed" early in life, may make individuals more susceptible to adverse health outcomes later in life when body reserves decline. Adverse influences in early life may alter gene expression and cell maturation or change the structure of affected organs. The effects of programming can also be mediated through changes in the homeostatic set-points of hormonal and metabolic systems.⁴ While research has focused primarily on fetal programming (Barker 1998), it has become increasingly clear that postnatal influences are also of utmost importance.

The factors that influence height may also influence cognitive development. A large literature indicates that height is correlated with intelligence. Although a portion of this correlation is driven by genes, environmental factors are also important (Lynn 1989). Sundet et al. (2005) used differences in cross-trait (height and intelligence), cross-twin correlations between monozygotic (MZ) and dizygotic (DZ) twin pairs to identify the roles played by shared environments and shared genes. They concluded that the environment plays a large role and is responsible for 65% of the height-intelligence correlation, with genes responsible for 35% of the observed correlation. These results are very similar to those on cross-trait, cross-twin correlations found much earlier by Husén (1959) in a large study of MZ and DZ twin pairs. Higher-birth-weight members of twin pairs are taller on average and have higher IQs than the lighter twins (Black et al. 2007). Other research using data on American twins has found that, within monozygotic twin pairs, greater fetal

4. References include Barker (1998), Bateson et al. (2004), Bertram and Hanson (2001), Dahlgren et al. (2001), Dickerson and Kemeny (2004), Drake and Walker (2004), Holmang (2001), Holness, Langdown, and Sugden (2000), Jackson (2000), Kajantie, Dunkel et al. (2003), Kajantie, Eriksson et al. (2003), Kajantie et al. (2004), Langley-Evans (2000), McMillen and Robinson (2005), Nilsson et al. (1998), Ozanne (2001), Phillips (2001), and Samuelsson et al. (2004).

growth is associated with greater stature and higher educational attainment (Behrman and Rosenzweig 2004).

The height-intelligence correlation may also be influenced by environmental conditions experienced in early childhood. Randomized experiments conducted in low-income environments have found that nutrition may be one such factor. In one study, nutritional supplements given to growth-retarded children improved their cognitive test scores, although these gains dissipated after the supplementation ended (Grantham-McGregor 2002; Walker et al. 2005). Research conducted in Guatemala found that children who had been provided with nutrition supplements during the first three years of life were taller, had higher scores on some cognitive tests, and (for girls) achieved higher educational attainment than children who did not receive supplements (Maluccio et al. 2009). Observational studies have also highlighted the role of early childhood. For example, Rääkkönen et al. (2009) used the Helsinki Birth Cohort Study to show that, conditional on health at birth, slower growth in height between birth and age 2 is associated with poorer performance on a range of intelligence tests in young adulthood.

Associations Between Height and Adult Outcomes Over the Life Course

The evidence indicates that height reflects a variety of environmental factors experienced *in utero* and in early childhood, and that the environmental factors that influence height may also affect health and cognitive development. Given this evidence, it is not surprising that height is associated with a wide range of economic and health outcomes in adulthood. Previous research indicates that height is associated with educational status, earnings, and occupational grade (Case and Paxson 2008a; Case, Paxson, and Islam 2008; Persico, Postlewaite, and Silverman 2004). Height is also associated with health and cognitive function at older ages (Case and Paxson 2008b). In this section, we summarize these associations using data from five American and British surveys that capture individuals at different points in the life course and that contain roughly comparable measures of adult outcomes.

The National Child Development Study (NCDS) tracks a cohort of all individuals born in Britain in a single week in 1958. The British Cohort Study (BCS) tracks a cohort of children born in a single week in 1970. For the NCDS, we use data from the surveys conducted when cohort members were 33 and 42 years of age; for the BCS, we use data from the survey conducted when the cohort members were 30 years old. The NCDS and BCS were designed to collect the same measures of health and economic status, providing comparable data on individuals in young to middle adulthood.

We also use data from the Panel Study of Income Dynamics (PSID), a longitudinal study of families, from which we draw information on household heads and their spouses between the ages of 25 and 55 who were interviewed between 1988 and 1997. Over this time period, the PSID collected detailed information on economic outcomes but relatively little on health.

In addition, we draw on the Whitehall II study, a longitudinal study of British civil servants. We use information collected from 1985 to 2001 on individuals between the ages of 34 and 71. Because individuals had to be employed in the British civil service when the study began, the sample is not representative of the population in Britain. (See Case and Paxson [2010] for details.) However, it contains a rich set of measures of health and cognitive functioning from middle age to early old age. Our final data source is the Health and Retirement Study (HRS), which is a panel study of Americans who were aged 51 to 61 years in 1992. We use information on individuals aged 55 and older who were interviewed between 1992 and 2004. The HRS contains information on employment and earnings (among those who were still employed) as well as health and cognitive function.

For each of these surveys, we select measures of economic and health status that are roughly comparable across the surveys, although not all measures were collected in all surveys. For the PSID and HRS, schooling is measured as reported years of education.

For the NCDS and the BCS, individuals are asked to report the age at which they left full-time schooling, and we measure years of education as the school-leaving age minus 5. Using this method, some individuals have implausibly high values for years of education, possibly because they resumed full-time schooling in adulthood after taking time off. For this reason, we truncate years of schooling at 20 for these two surveys.⁵ We also analyze employment and the logarithm of average hourly earnings (conditional on employment). Our health measures include self-reported health status, activity limitations and disability, problems with activities of daily living, depression, and cognitive function. More details on these measures are provided below. We regress these measures of economic and health outcomes on height in inches plus, where appropriate, indicators for age, ethnicity, sex, and survey wave. In cases where there are multiple observations for an individual at different time periods, we use clustered standard errors.

Table 1 reports results for the economic outcomes. Results for men and women are reported separately because of their different rates of labor force participation. We do not include results from the Whitehall study in this table. Although the data have a rich set of health measures, they do not include measures of hourly earnings, and there are numerous missing values for the measure of the school-leaving age. Furthermore, selection into the sample was, by construction, contingent on being employed at the start of the study; thus, the association between height and employment is of less interest. For the HRS, we use only individuals ages 55 to 74 because labor force participation is very low at older ages.

The results in Table 1 indicate that height is associated with economic outcomes in all the samples. Each inch of height is associated with an increase in schooling of between 0.05 years (for 30-year-old women in the BCS) and 0.16 years (for men in the HRS).⁶ All coefficients on height in the schooling regressions are statistically different from zero. Although we do not present results from the Whitehall study here, in other work, we have shown that height is significantly associated with the highest level of educational qualification received (Case and Paxson 2010). Employment is also associated with height for most groups. For men aged 55 and under, each inch of height is associated with an increase in the probability of working of between 0.2 and 0.6 percentage points. Women have similar coefficients, although these are less precisely estimated among women in the NCDS. All groups show a substantial height premium in average hourly earnings, ranging from 0.012 to 0.028 for men and 0.007 to 0.027 for women. The magnitudes of these coefficients are economically significant. For example, a height premium of 0.025 for men in the NCDS and the HRS implies that a 4-inch increase in height (a move from the 25th to 75th percentile in height) is associated with a 10% increase in hourly earnings.

Table 2 presents evidence on the association between height and a variety of health and cognitive outcomes in adulthood. All surveys ask respondents to rate their health. For the NCDS and BCS, self-reported health is measured on a four-point scale (1 = excellent, 2 = good, 3 = fair, and 4 = poor). The other three studies use five-point scales (1 = excellent, 2 = very good, 3 = good, 4 = fair, and 5 = poor). The NCDS and BCS ask questions about whether the individual has a long-standing illness or is registered as disabled; comparable questions do not exist in the other surveys. Whitehall II and HRS, both of which surveyed

5. The NCDS, BCS, and Whitehall II studies collected detailed information on highest academic qualification attained by cohort members. These are better measures of educational achievement, and ordered probits reveal a significant association between highest degree and height in all three studies. Case et al. (2008) found, for the educational attainment of men followed in the British Household Panel Study, that each inch of height has the same effect on the ordered probit index as a rollback of the age clock by five years, so that movement from the 25th to the 75th percentile in the height distribution has an association with educational attainment comparable to moving to a younger (and more highly educated) generation.

6. The association between height and completed education appears to be larger in the HRS than in the surveys used here. It is not clear why this would be the case, but it may reflect differences in the construction of education variables between the U.S. and the U.K. surveys or greater selection by height into education in the (older) HRS cohorts.

Table 1. Regressions of Education, Employment, and Earnings on Height in Inches

Dependent Variable	NCDS, Age 33	NCDS, Age 42	BCS, Age 30	PSID, Ages 25–55	HRS, Ages 55–74
Men					
Schooling (years)	0.0913 (0.0114)		0.0679 (0.0104)	0.0686 (0.0111)	0.1571 (0.0164)
Employed (indicator)	0.0061 (0.0015)	0.0041 (0.0015)	0.0037 (0.0014)	0.0018 (0.0008)	0.0015 (0.0019)
Log average hourly earnings, if employed	0.0262 (0.0037)	0.0248 (0.0042)	0.0122 (0.0023)	0.0145 (0.0029)	0.0277 (0.0051)
Maximum observations	5,833		5,424	31,996	27,606
Women					
Schooling (years)	0.0940 (0.0107)		0.0550 (0.0104)	0.0454 (0.0106)	0.1185 (0.0119)
Employed (indicator)	0.0036 (0.0025)	0.0028 (0.0022)	0.0093 (0.0021)	0.0023 (0.0010)	0.0029 (0.0017)
Log average hourly earnings, if employed	0.0170 (0.0042)	0.0195 (0.0048)	0.0114 (0.0025)	0.0067 (0.0030)	0.0272 (0.0044)
Maximum observations	5,815		5,757	31,999	35,927

Notes: Each cell contains a coefficient and standard error on height in inches, from a regression of the dependent variable on height and a set of ethnicity indicators. For results using the PSID and HRS, indicators for year of age are also included. For all outcomes except schooling, the PSID and HRS contain multiple observations per person; thus, survey wave/year dummy variables are included, and standard errors are clustered at the individual level. “Schooling” is defined as reported years of schooling for the PSID and HRS, and as the age at which the individual left full-time education minus 5 for the NCDS and BCS. In some cases, regressions contain smaller numbers of observations due to missing values of the dependent variables.

older individuals, contain information on problems with activities of daily living (ADLs). For Whitehall II, our measure of ADLs is the number of problems the individual has with seven items: problems with vigorous activity, bathing or dressing, moderate activity, lifting, bending, walking several miles, and climbing several flights of stairs. For the HRS, we use six activities that reflect serious limitations: problems with dressing, walking across the room, bathing, eating, getting in and out of bed, and using the toilet. Both the HRS and Whitehall II surveys contain a measure of depression. Finally, both Whitehall II (in Waves 3 and 5) and the HRS (in Waves 3 through 6) conducted cognitive tests. For Whitehall II, we show results for tests of word recall and ability in arithmetic. For the HRS, we use a test of delayed word recall and an indicator of whether the individual could count backward from 10.

In all samples, taller individuals report themselves to be in better health (remembering that a lower number refers to a better health assessment). In the NCDS and BCS, taller individuals are also less likely to report having a long-standing illness or to be registered as disabled. The magnitudes of these coefficients are large, in light of the fact that relatively few of these younger adults have serious illness or are disabled. For example, in the BCS, 23.2% of individuals report a long-standing illness, and 1.5% are registered as disabled. The results imply that a 4-inch increase in height would reduce the probability of a long-standing illness from 23.2% to 21.4% and reduce the probability of disability from 1.5% to 0.9%.

The association between height and health persists into older adulthood. Results from Whitehall II and the HRS indicate that taller individuals have fewer problems with activities of daily living and are less likely to be depressed. Although the two surveys use different measures of depression, the results for both imply that a 4-inch increase in height

Table 2. Regressions of Measures of Health and Cognition on Height in Inches

Dependent Variable	NCDS, Age 33	NCDS, Age 42	BCS, Age 30	PSID, Ages 25–55	Whitehall II, Ages 34–71	HRS, Ages 55+
Self-reported Health Status ^a	-0.0171 (0.0026)	-0.0150 (0.0028)	-0.0126 (0.0023)	-0.0085 (0.0027)	-0.0241 (0.0032)	-0.0118 (0.0031)
Has Long-standing Illness	-0.0010 (0.0014)	-0.0036 (0.0017)	-0.0044 (0.0014)			
Is Registered as Disabled	-0.0023 (0.0004)	-0.0016 (0.0006)	-0.0016 (0.0004)			
Number of ADLs ^b					-0.0141 (0.0004)	-0.0024 (0.0020)
Depression Score ^c					-0.0758 (0.0260)	-0.0214 (0.0047)
Word Recall Score ^d					0.0574 (0.0101)	0.0434 (0.0046)
Counting Backward ^c						0.0041 (0.0007)
Numerical Ability Score ^f					0.4419 (0.0278)	
Maximum Observations	11,648		11,181	63,995	29,774	66,269

Notes: Each cell contains a coefficient and standard error on height in inches, from a regression of the dependent variable on height, an indicator for the respondent's sex, and (for all but the Whitehall II survey) a set of ethnicity indicators. For the PSID, Whitehall, and HRS, indicators for year of age are included. The PSID, Whitehall, and HRS contain multiple observations per person; thus, survey wave/year indicators are included, and standard errors are clustered at the individual level. For some outcomes, regressions contained smaller numbers of observations because data were not collected in all waves, or because of missing values of the dependent variables.

^aFor the NCDS and BCS, 1 = excellent, 2 = good, 3 = fair, and 4 = poor; for the PSID, Whitehall, and HRS, 1 = excellent, 2 = very good, 3 = good, 4 = fair, and 5 = poor.

^bFor Whitehall, range = 0–7, mean = 0.28; for the HRS, range = 0–5, mean = 0.26.

^cFor Whitehall, range = 0–45, mean = 13.8, standard deviation = 7.3; for the HRS, range = 0–8, mean = 1.4, standard deviation = 1.9.

^dFor Whitehall, range = 0–18, mean = 6.4, standard deviation = 2.4; for the HRS, range = 0–10, mean = 4.6, standard deviation = 2.1;

^eCoded as equal to 1 if the individual could accurately count backward from 10, and 0 if not. Mean = 0.88.

^fRange = 0–33, mean = 22.4, standard deviation = 6.5.

is associated with a decline in the depression score of 4% of 1 standard deviation of the depression measure. Taller individuals also perform significantly better on cognitive tests, even among the (younger) Whitehall II sample.

Interpretation of Results

The finding that height is associated with a wide range of economic and health outcomes throughout adulthood provides support for the hypothesis that early childhood circumstances have long-lasting effects. However, as discussed in the conceptual framework, there are several routes through which childhood circumstances could influence adult outcomes. One possibility is that height operates solely through educational attainment and economic status in early adulthood: children who are healthier and are raised in more advantaged circumstances achieve higher levels of education and get better first jobs, which in turn improves economic opportunities and health trajectories over adulthood. If so, then the “separability” discussed above would hold, and information on early adulthood would be sufficient to

capture the effects of early-life circumstances. Another possibility is that, although childhood circumstances may influence education and economic status in early adulthood, they may also exert independent effects on long-run health and economic outcomes.

A number of studies have examined this issue by estimating reduced-form models similar to those presented above (with additional controls for child health and economic circumstances, if available) and then examining whether the addition of controls for characteristics in young adulthood account for the association between childhood circumstances and later-life outcomes. In some cases, the evidence suggests that childhood health influences adult outcomes, even controlling for outcomes in young adulthood. For example, Case et al. (2005) showed, after they controlled for measures of socioeconomic status in childhood, that chronic conditions in childhood and height at age 16 predict socioeconomic status of 42-year-old men from the NCDS. This predictive power of chronic conditions and height at age 16 for socioeconomic status at age 42 continues even after education, health, and socioeconomic status at ages 23 and 33 are controlled for. The same is true when examining self-reported health status at age 42.

Case and Paxson (2008b) found different results when using the HRS to examine the effects of height on a set of cognitive and health measures among older adults. In this case, although the reduced-form estimates show that height predicts cognition and health, the coefficients on height become much smaller and are typically not significantly different from zero when controls for education are added. Older individuals who are more highly educated have better cognitive function and health status. It is unknown whether the “protective” effects of education are causal or whether greater education reflects better cognitive ability and health in childhood.

To extend this research, we reestimated the models shown in Tables 1 and 2 and included controls for schooling, which are available for all of the data sets we use. (For Whitehall II, for which we do not have years of education, we included indicator variables for the highest degree attained.) In all cases, schooling is significantly associated with economic and health outcomes in adulthood. However, although the coefficients on height decline when controls for schooling are added, height continues to be significantly associated with adult outcomes in most cases. For the BCS, which assesses outcomes at the earliest stage of adulthood, the addition of schooling causes the coefficients on height to decline between 5% (for whether disabled) to 28% (for hourly earnings for women). The median decline in the coefficient on height is 30.3% for the NCDS, 54.9% for the PSID, and 7.5% for Whitehall II. Consistent with Case and Paxson (2008b), the associations between height and outcomes in old age in the HRS become much smaller once controls for years of schooling are included (the median decline is 79.2%). The smallest declines among the HRS results are for average hourly earnings and the two cognitive tests, which remain significant even after schooling is included.

These results indicate that height, as a marker of early childhood circumstances, is associated with health, cognitive, and economic outcomes throughout adulthood. This association appears to be only partially mediated by schooling. Although these results underscore the importance of early childhood environment, they provide little information about what specific aspects of early childhood are being captured by height. We turn to this issue in the following section, where we use data from the NLSY79 Children and Young Adults survey⁷ to document the associations between height, health, and cognitive development among children.

7. The NLSY79 is a nationally representative longitudinal study of approximately 12,000 young men and women who were ages 14 to 22 when the study began in 1979. A complete description of these data is available online at <http://www.bls.gov/nls/nlsy79ch.htm>.

HEIGHT AND EARLY-LIFE HEALTH: EVIDENCE FROM THE NLSY

Beginning in 1986, data have been collected on the children of original female NLSY79 cohort members as part of the NLSY Child and Young Adults survey. These data provide information on a rich set of outcomes, including prenatal care and health at birth, children's cognitive function, schooling, and self-perceptions of scholastic competence and self worth. In addition, all children born to an original NLSY79 woman can be identified as maternal siblings, which allows us to look at the associations between heights and developmental outcomes while controlling for maternal fixed effects. We use data on children who are 14.5 years of age and younger, collected every other year from 1986 to 2006.⁸

Table 3 provides descriptive statistics for these children. Eighteen percent of them were born to teen mothers, while 8% were born to mothers older than age 35. Approximately 8% were born at low birth weight (<2,500 grams). One in three was born to a mother who smoked during pregnancy, and one in five was born to a mother who did not seek prenatal care in the first trimester.

We have information on children's heights and cognitive test score results, both of which were collected in multiple waves of the survey. Information on mothers includes her educational attainment, whether she lived with both parents at age 14, whether her family received magazines and newspapers when she was 14, whether her family had a library card when she was 14, the educational attainment of her parents, her height, and her ethnicity. We will include information on mother's characteristics in some regression specifications and mother fixed effects in others to investigate whether there are unobservable mother-specific characteristics that affect children's heights and their development.

Table 4 presents evidence on the association between height and cognitive ability in childhood. The first panel presents results on the Peabody Individual Achievement Test (PIAT) math and word recognition scores for children at ages 5 and 6. The second panel presents results for children at ages 10 and 11. In addition to the earlier tests, older children are tested on PIAT reading comprehension, digit span, and the Peabody Picture Vocabulary Test (PPVT). The bottom panel includes test score results for children at all ages. Each coefficient presented in Table 4 is the coefficient on the child's height-for-age z score in regressions of cognitive tests that include indicators for the child's age at testing (rounded to the nearest 6 months), an indicator for the child's sex, whether he or she was born to a teenaged mother or an older mother (older than 35), the child's birth order, and a set of race and survey year indicators.

For all tests at all ages, we find a positive and significant association between test scores and heights in ordinary least squares (OLS) regressions, with associations becoming larger at older ages. With two exceptions, we continue to find a positive and significant association when mother fixed effects are added to regressions. The addition of mother fixed effects diminishes the association between height and cognitive tests for most tests, but does not eliminate it. The coefficient on height, in regressions that include mother fixed effects, is identified solely from differences in test scores between siblings, when tested at a given age, and differences in their height-for-age z scores at that age.

Siblings could differ in their heights and their test scores because the household environment one was born into could have been healthier, an explanation we will examine later in this section. Differences in heights could also be due to differences in the genetic material inherited from parents or to the interaction between genes and environment. Without more information, it is not possible to say more than this. However, we can say that even after we control for the genetic material that siblings share and for the part of their

8. Older children are included in the survey. However, because the NLSY79 consists of women born between 1958 and 1965, the older children in the sample were born to an increasingly selected sample of younger mothers.

Table 3. Descriptive Statistics, NLSY Child and Young Adult Sample (ages 0–14.5 years)

Characteristic	Mean	SD	Number of Observations
Child Characteristics (one observation per child)			
Mother was teenager at child's birth	0.181		10,324
Mother was older than 35 at child's birth	0.078		10,324
Child's birth order	1.97		10,324
Child female	0.489		10,324
Child born at low weight	0.075		10,318
Mother smoked during pregnancy	0.322		9,422
Mother had no prenatal care	0.015		9,200
Mother began prenatal care in first trimester	0.799		9,200
Mother began prenatal care in second trimester	0.144		9,200
Mother began prenatal care in third trimester	0.039		9,200
Child Characteristics (multiple observations per child)			
PIAT math test	100.35	14.08	31,175
PIAT reading recognition test	103.95	14.85	31,045
PIAT reading comprehension test	101.17	13.96	26,199
Digit span test (range = 1–19)	9.78	3.12	16,968
Peabody Picture Vocabulary Test (PPVT)	90.45	20.30	17,341
Height-for-age <i>z</i> score	0.137	1.36	46,671
Mother Characteristics			
Highest grade attained	13.21		3,778
Lived with intact family at age 14	0.745		4,433
Family at age 14 received magazines	0.524		4,445
Family at age 14 received newspaper	0.724		4,447
Family at age 14 had library card	0.695		4,446
Highest grade attained by mother's mother	10.56		4,206
Highest grade attained by mother's father	10.59		3,770
Mother's adult height (inches)	64.24		3,248
Mother Hispanic	0.179		4,471
Mother black	0.267		4,471

home environment that is constant, children who are taller outperform shorter siblings on cognitive tests throughout childhood.

We also examine whether taller children progress more rapidly through school. We use two measures of progress in school: the highest grade attained (top panel) and an indicator of whether a child is at the appropriate grade given his or her age (bottom panel). OLS regression coefficients on height are presented in columns 1 and 3 of Table 5, and those from regressions that include mother fixed effects are presented in columns 2 and 4. Results are reported for children ages 6–14. Height-for-age *z* scores at age 5 are significant predictors of children's progress through school, with or without controls for mother fixed effects. Using information on height at age 5 interacted with age (columns 3 and 4), we test whether this

Table 4. Heights and Test Scores

	PIAT Math	PIAT Recognition	PIAT Comprehension	Digit Span	PPVT
Regressions of Test Scores on Height-for-Age <i>z</i> Scores, 5- and 6-Year-Olds					
No mother fixed effects	0.637 (0.124)	0.395 (0.121)			
Mother fixed effects	0.402 (0.183)	0.267 (0.180)			
Number of observations	6,492	6,384			
Regressions of Test Scores on Height-for-Age <i>z</i> Scores, 10- and 11-Year-Olds					
No mother fixed effects	0.734 (0.151)	0.559 (0.161)	0.548 (0.147)	0.133 (0.035)	0.753 (0.196)
Mother fixed effects	0.744 (0.218)	0.299 (0.221)	0.463 (0.208)	0.091 (0.052)	0.201 (0.252)
Number of observations	6,206	6,192	6,133	6,162	6,105
Regressions of Test Scores on Height-for-Age <i>z</i> Scores, All Ages					
No mother fixed effects	0.713 (0.087)	0.587 (0.095)	0.520 (0.087)	0.175 (0.024)	0.971 (0.136)
No mother fixed effects, mother controls ^a	0.642 (0.108)	0.573 (0.116)	0.535 (0.108)	0.167 (0.030)	0.925 (0.167)
Mother fixed effects	0.376 (0.078)	0.192 (0.083)	0.248 (0.084)	0.089 (0.025)	0.362 (0.148)
Number of observations	30,421	30,296	25,606	16,579	16,853

Notes: All models include indicators for the child's age (rounded to the nearest 6 months) at testing, an indicator for the child's sex, whether he or she was born to a teen-aged mother or an older mother (older than 35), the child's birth order, and a set of indicators for race and survey year. The models with "mother controls" include the mother's highest grade; indicators for whether the mother (at age 14) lived in a family that was intact, received magazines, received a newspaper, had a library card, and lived in an urban area; the highest grade attained by her mother and father; and her adult height. For the top two panels, there is one observation per child. In the bottom panel, with multiple observations per child, the standard errors are clustered at the child level.

^aSamples with mother controls included are smaller than those noted in table because of missing values.

is because taller children start school earlier or because they advance more quickly through grades. At age 6, taller siblings are no more likely than shorter siblings to be at appropriate grade-for-age: results including mother fixed effects suggest that a taller and shorter sibling differing by 1 standard deviation in height-for-age *z* scores are, at age 6, equally likely to reach grade-for-age ($-0.0179 + 0.0024 \times 6 = -0.0035$). However, by age 14, the taller sibling is 1.6 percentage points more likely to be observed at appropriate grade-for-age.

There are several reasons why taller children may progress more rapidly through school. One is that they have better cognitive skills. Another is that taller children have greater self-esteem, perhaps because they are treated differently by parents, teachers, or peers. We can distinguish between these two possibilities by using data that were collected directly from the children in the NLSY Child and Young Adult Survey. At ages 12–14, children were asked a battery of questions about how well they perform at school and how pleased they are with their lives. Each individual item is scored from 1 (low) to 4 (high), and the Total Scholastic Competence score and Total Self-Worth score are sums of

Table 5. Grade Level in School and Height: Children Ages 6–14

	(1)	(2)	(3)	(4)
Highest Grade Attained				
Height-for-age <i>z</i> score at age 5	0.0177 (0.004)	0.0086 (0.0051)	-0.0145 (0.0095)	-0.0287 (0.0109)
Height-for-age <i>z</i> score at age 5 × age			0.0033 (0.0010)	0.0038 (0.0011)
At Appropriate Grade Level at a Given Age (indicator, mean = 0.873)				
Height-for-age <i>z</i> score at age 5	0.0118 (0.0023)	0.0056 (0.0029)	-0.0092 (0.0048)	-0.0179 (0.0061)
Height-for-age <i>z</i> score at age 5 × age			0.0022 (0.0006)	0.0024 (0.0007)
Mother fixed effects?	No	Yes	No	Yes

Notes: Number of observations = 23,783. “At appropriate grade level” is defined to equal 1 if the children over 7 years and 6 months of age have completed grade 1, if children over 8 years and 6 months have completed grade 2, and so on. All models include a set of indicators for the child’s age rounded to the nearest 6 months, an indicator for the child’s sex, whether he or she was born to a teen-aged mother or an older mother (older than 35), the child’s birth order, and a set of race and survey year indicators. Standard errors are clustered at the child level.

individual items. Column 2 of Table 6 presents regression results in which a scholastic or self-worth score is regressed on height-for-age *z* scores, with controls for mother and child characteristics. Results in column 3 control in addition for mothers’ fixed effects.

We find that taller children—and, within sibships, taller siblings—are more likely to report that they are good students. On average, their Total Scholastic Competence score is significantly higher, with or without mother fixed effects. These results are consistent with evidence that taller children have better cognitive test scores and progress through school more quickly than less tall children and siblings. In contrast, taller children and taller siblings are not more likely to report being pleased with themselves or that they are happy with how they do things. Taller children are significantly *less* likely to report that they are “happy with the way I am.” Total Self-Worth scores are not significantly associated with height. The associations we find between height and cognitive test scores, educational attainment, and self-perceptions of scholastic competence do not appear to spill over into children’s sense of their worth.

Previous research has indicated that physical growth is related to children’s health. Results from the NLSY79 Child and Young Adults survey are consistent with this evidence. Children’s heights are significantly associated with their health in childhood. Table 7 presents evidence that taller children, and taller siblings within sibships, are significantly less likely to have a limiting condition. Most prominently, taller children are less likely to be reported as having an emotional or neurological limiting condition.

Finally, the Children and Young Adults data allow us to investigate why some children, and within a sibship why some siblings, are taller than others. Table 8 shows the association between height-for-age *z* scores in childhood and children’s birth weight, birth length, and mothers’ prenatal health. We find that, with or without mother fixed effects, children who were heavier and longer at birth are significantly taller throughout childhood. This suggests that even within a sibship, it may be difficult to place smaller newborns onto the same trajectory that larger siblings are on. Controlling for birth weight and length, we find that children born to mothers who received no prenatal care have height-for-age *z* scores that are two-tenths of a standard deviation shorter on average. This is true for comparisons across all children in the OLS regression (column 1) and in mother fixed effect regressions,

Table 6. Child's Self-perception and Height: NSLY Child and Young Adult Survey, Ages 12–14

	Mean of Dependent Variable	Coefficient on Height-for-Age z Score	
		No Fixed Effects	Mother Fixed Effects
Scholastic Competence			
Good at school work	2.93	0.030 (0.011)	0.025 (0.018)
As smart as other kids the same age	3.06	0.035 (0.011)	0.019 (0.018)
Does schoolwork quickly	2.84	0.042 (0.012)	0.041 (0.020)
Remembers things easily	2.78	0.034 (0.011)	0.039 (0.018)
Does class work well	3.22	0.013 (0.010)	0.012 (0.017)
Can figure out answers at school	2.66	0.028 (0.011)	0.027 (0.019)
Total Scholastic Competence score	17.50	0.183 (0.046)	0.164 (0.072)
Self-Worth			
Pleased with self	3.44	-0.010 (0.009)	0.008 (0.014)
Like the way leading life	3.41	-0.013 (0.010)	-0.004 (0.016)
Happy with self as a person	3.60	-0.008 (0.008)	-0.002 (0.013)
Like who I am	3.41	0.008 (0.009)	0.013 (0.016)
Happy with the way I am	3.43	-0.020 (0.009)	-0.029 (0.015)
Happy with how I do things	3.29	0.004 (0.010)	0.010 (0.017)
Total Self-Worth score	20.58	-0.043 (0.037)	-0.011 (0.057)

Notes: Number of observations = 7,438. Each individual item is scored from 1 (low) to 4 (high), and the total scores are sums of the individual items. All models include a set of indicators for the child's age rounded to the nearest 6 months, an indicator for the child's sex, whether he or she was born to a teen-aged mother or an older mother (older than 35), the child's birth order, and a set of race and survey year indicators. Standard errors are clustered at the child level.

where identification comes from comparing mothers' behaviors and children's heights within sibships (column 2).

We find that maternal smoking and drinking while pregnant, in the presence of controls for birth length and weight, do not appear to affect height in childhood. However, this is because their effects are largely realized in birth weight and length. Columns 5 and 6 of Table 8 report fixed-effect results on the associations between birth weight, birth length, and mothers' behaviors while pregnant. Controlling for all the characteristics of mothers that do not change between pregnancies, we find birth weight is significantly and negatively

Table 7. Regressions of Health Measures on Height-for-Age z Scores

	Has Limiting Condition	Has a Limiting Condition That Is:		
		Emotional/ Neurological	Respiratory	Other
Mean of Dependent Variable	0.100	0.022	0.045	0.040
No Mother Fixed Effects	-0.0050 (0.0014)	-0.0026 (0.0007)	0.0004 (0.0009)	-0.0031 (0.0009)
Mother Fixed Effects	-0.0031 (0.0013)	-0.0017 (0.0007)	0.0007 (0.0008)	-0.0018 (0.0009)

Notes: Number of observations = 46,101. All models include indicators for the child's age (rounded to the nearest 6 months) at testing, an indicator for the child's sex, whether he or she was born to a teen-aged mother or an older mother (older than 35), the child's birth order, and a set of race and survey year indicators. Standard errors are clustered at the child level.

Table 8. Prenatal Health and Height

Variable	Height-for-Age z Score in Childhood				Birth Weight z Score	Birth Length z Score
	Birth Weight z Score	0.147 (0.013)	0.105 (0.015)			
Birth Length z Score	0.098 (0.010)	0.093 (0.011)				
No Prenatal Care	-0.213 (0.101)	-0.227 (0.107)	-0.266 (0.106)	-0.264 (0.111)	-0.028 (0.127)	-0.325 (0.181)
Prenatal Care, Second Trimester	0.026 (0.033)	0.047 (0.033)	0.005 (0.034)	0.026 (0.034)	-0.130 (0.038)	-0.051 (0.055)
Prenatal Care, Third Trimester	0.082 (0.055)	0.096 (0.051)	0.085 (0.057)	0.095 (0.052)	-0.057 (0.066)	0.064 (0.095)
Smoking Frequency	-0.008 (0.018)	-0.022 (0.032)	-0.067 (0.019)	-0.040 (0.033)	-0.097 (0.035)	-0.071 (0.050)
Drinking Frequency	0.009 (0.010)	0.004 (0.012)	0.004 (0.010)	-0.003 (0.123)	-0.023 (0.014)	-0.060 (0.020)
Mother Fixed Effects?	No	Yes	No	Yes	Yes	Yes
Number of Observations	37,568				8,088	8,088

Notes: Regressions for which the dependent variable is the height-for-age z score include indicators for the child's age (rounded to the nearest 6 months), an indicator for the child's sex, whether he or she was born to a teen-aged mother or an older mother (older than 35), the child's birth order, and a set of race and survey year indicators. Standard errors are clustered at the child level. Regressions for which the dependent variable is birth weight or birth length include a cubic for the mother's age at the time of the birth, an indicator for the child's sex, whether he or she was born to a teen-aged mother or an older mother (older than 35), the child's birth order, and a set of race and survey year indicators. Standard errors are clustered at the child level.

associated with smoking during pregnancy, while birth length is significantly and negatively associated with reporting no prenatal care and with drinking alcohol while pregnant.

CONCLUSION

There are several reasons why it is important to study the long-run consequences of health in childhood. Knowledge about the effects of childhood health problems on economic and health outcomes later in life is useful for assessing the benefits of interventions that improve child health. Furthermore, an understanding of the childhood antecedents of adult

disease may be useful for making projections of future trends in adult morbidity and disability. However, without data that follow individuals from childhood to adulthood, it is difficult to quantify the consequences of childhood health. We have shown how height, which reflects (in part) environmental conditions in the prenatal and early childhood periods, can be used to understand the long-run consequences of childhood health.

Data from the NLSY79 Children and Young Adults survey demonstrate that within families, a mother's behavior during pregnancy can result in significant differences between her children—differences that may play out over the life course. Among children in general, and more strikingly among siblings, those who are heavier and longer at birth are, on average, taller in childhood. Among children in general, and among siblings more specifically, children who are taller are healthier, score significantly higher on cognitive tests and make their way more quickly through school. They are also more likely to perceive themselves to be scholastically competent. All of these characteristics send children on different trajectories into adulthood and old age—trajectories that were, in part, set in place by mothers' behaviors while pregnant.

The evidence presented here on the associations between height and outcomes throughout adulthood indicate that childhood circumstances are important. Taller individuals attain more education, earn more, and are more likely to be employed. They are also more likely to have better health and cognitive outcomes in middle and older ages. A substantial portion of the better outcomes of taller individuals appears to be due to their higher levels of educational attainment.

When considering these results, it is important to keep in mind that the environments in which children are being raised today are substantially different from those in which the oldest individuals we examine—members of the HRS and Whitehall II—were raised as children. These older individuals grew up in environments in which major causes of child deaths included diarrhea, pneumonia, and diseases such as diphtheria, measles, and typhoid (Case and Paxson 2009). A large body of literature indicates that exposures to infectious diseases can have adverse implications for cognitive development and adult health (Costa 2000; Finch and Crimmins 2004). It may be tempting to speculate that with the decline in infectious disease, childhood health will have a smaller role in determining adult outcomes. However, the fact that less-healthy children born in the last two decades are shorter, and that these shorter children have worse performance on cognitive tests and slower progress through school, indicates that child health is likely to remain an important determinant of health and economic status once these children reach adulthood.

REFERENCES

- Adler, N.E., T. Boyce, M.A. Chesney, S. Cohen, S. Folkman, R.L. Kahn, and S.L. Syme. 1994. "Socioeconomic Status and Health: The Challenge of the Gradient." *American Psychologist* 49:15–24.
- Almond, D. 2006. "Is the 1918 Influenza Pandemic Over? Long-Term Effects of *in Utero* Influenza Exposure in the Post-1940 U.S. Population." *Journal of Political Economy* 114:672–712.
- Barker, D.J.P. 1992. "The Fetal Origins of Diseases of Old Age." *European Journal of Clinical Nutrition* 46 (Suppl. 3):S3–S9.
- . 1995. "Fetal Origins of Coronary Heart Disease." *British Medical Journal* 311:171–74.
- . 1998. "Programming the Baby." Pp. 13–41 in *Mothers, Babies and Health in Later Life*, 2nd ed. Edinburgh: Churchill Livingstone.
- Bateson, P., D.J.P. Barker, T. Clutton-Brock, D. Deb, B. D'Udine, R.A. Foley, P. Gluckman, K. Godfrey, T. Kirkwood, M.M. Lahr, J. McNamara, N.B. Metcalfe, P. Monaghan, H.G. Spencer, and S.E. Sultan. 2004. "Developmental Plasticity and Human Health." *Nature* 430:419–21.
- Beard, A.S. and M.J. Blaser. 2002. "The Ecology of Height: The Effect of Microbial Transmission on Human Height." *Perspectives in Biology and Medicine* 45: 475–99.

- Behrman, J.R. and M.R. Rosenzweig. 2004. "Returns to Birthweight." *Review of Economics and Statistics* 86:586–601.
- Bertram, C.E. and M.A. Hanson. 2001. "Animal Models and Programming of the Metabolic Syndrome." *British Medical Bulletin* 60:103–21.
- Black, S.E., P.J. Devereux, and K.G. Salvanes. 2007. "From the Cradle to the Labor Market? The Effect of Birth Weight on Adult Outcomes." *Quarterly Journal of Economics* 122: 409–39.
- Brunner, E., M.J. Shipley, D. Blane, G. Davey Smith, and M.G. Marmot. 1999. "When Does Cardiovascular Risk Start? Past and Present Socioeconomic Circumstances and Risk Factors in Adulthood." *Journal of Epidemiology and Community Health* 53:757–64.
- Case, A., A. Fertig, and C. Paxson. 2005. "The Lasting Impact of Childhood Health and Circumstance." *Journal of Health Economics* 24:365–89.
- Case, A. and C. Paxson. 2008a. "Stature and Status: Height, Ability and Labor Market Outcomes." *Journal of Political Economy* 116:499–532.
- . 2008b. "Height, Health and Cognitive Function at Older Ages." *American Economic Review Papers and Proceedings* 98:463–67.
- . 2009. "Early Life Health and Cognitive Function in Old Age." *American Economic Review Papers and Proceedings* 99:104–109.
- . 2010. "The Long Reach of Childhood Health and Circumstance: Evidence From the Whitehall II Study." NBER Working Paper No. 15640. National Bureau of Economic Research, Cambridge, MA.
- Case, A., C. Paxson, and M. Islam. 2008. "Making Sense of the Labor Market Height Premium: Evidence From the British Household Panel Study." *Economic Letters* 102:174–76.
- Chandola, T., M. Bartley, A. Sacker, C. Jenkinson, and M. Marmot. 2003. "Health Selection in the Whitehall II Study, UK." *Social Science and Medicine* 56:2059–72.
- Costa, D. 2000. Causes of Improving Health and Longevity at Older Ages: A Review of the Explanations." *Genus* LXI(1):21–38.
- Crimmins, E.M. and C.E. Finch. 2006. "Infection, Inflammation, Height and Longevity." *Proceedings of the National Academy of Sciences* 103:498–503.
- Currie, J. 2009. "Healthy, Wealthy and Wise: Socioeconomic Status, Poor Health in Childhood, and Human Capital Development." *Journal of Economic Literature* 47:87–122.
- Cutler, D. and A. Lleras-Muney. 2008. "Education and Health: Evaluating Theories and Evidence." Chapter 2 in *Making American Healthier*, edited by R.F. Schoeni, J. House, G.A. Kaplan, and H. Pollack. New York: Russell Sage Foundation.
- Dahlgren, J., C. Nilsson, E. Jennische, H.P. Ho, E. Eriksson, A. Niklasson, P. Bjorntorp, K. Albertsson Wikland, and A. Holmang. 2001. "Prenatal Cytokine Exposure Results in Obesity and Gender-Specific Programming." *American Journal of Physiology–Endocrinology and Metabolism* 281:E326–E34.
- Dickerson, S.S. and M.E. Kemeny. 2004. "Acute Stressors and Cortisol Responses: A Theoretical Integration and Synthesis of Laboratory Research." *Psychological Bulletin* 130:355–91.
- Drake, A.J. and B.R. Walker. 2004. "The Intergenerational Effects of Fetal Programming: Non-genomic Mechanisms for the Inheritance of Low Birth Weight and Cardiovascular Risk." *Journal of Endocrinology* 180:1–16.
- Finch, C.E. and E.M. Crimmins. 2004. "Inflammatory Exposures and Historical Changes in Human Life Spans." *Science* 305:1736–39.
- Grantham-McGregor, S. 2002. "Linear Growth Retardation and Cognition." *Lancet* 359:542.
- Holmang, A. 2001. "Perinatal Origin of Adult Disease." *Scandinavian Cardiovascular Journal* 35:178–85.
- Holness, M.J., M.L. Langdown, and M.C. Sugden. 2000. "Early-Life Programming of Susceptibility to Dysregulation of Glucose Metabolism and the Development of Type 2 Diabetes Mellitus." *Biochemical Journal* 349:657–65.
- Husén, T. 1959. *Psychological Twin Research*. New York: Free Press.

- Jackson, A.A. 2000. "Nutrients, Growth, and the Development of Programmed Metabolic Function." *Advances in Experimental Medicine and Biology* 478:41–55.
- Kajantie, E., L. Dunkel, U. Turpeinen, U.H. Stenman, P.J. Wood, M. Nuutila, and S. Andersson. 2003. "Placental 11 β -Hydroxysteroid Dehydrogenase-2 and Fetal Cortisol/Cortisone Shuttle in Small Preterm Infants." *Journal of Clinical Endocrinology and Metabolism* 88:493–500.
- Kajantie, E., J. Eriksson, D.J.P. Barker, T. Forsén, C. Osmond, P.J. Wood, S. Andersson, L. Dunkel, and D.I.W. Phillips. 2003. "Birthsize, Gestational Age and Adrenal Function in Adult Life: Studies of Dexamethasone Suppression and ACTH₁₋₂₄ Stimulation." *European Journal of Endocrinology* 149:569–75.
- Kajantie, E., J. Eriksson, C. Osmond, P.J. Wood, T. Forsén, D.J.P. Barker, and D.I. Phillips. 2004. "Size at Birth, the Metabolic Syndrome and 24-h Salivary Cortisol Profile." *Clinical Endocrinology* 60:201–207.
- Langley-Evans, S.C. 2000. "Critical Differences Between Two Low Protein Diet Protocols in the Programming of Hypertension in the Rat." *International Journal of Food Sciences and Nutrition* 51:11–17.
- Li, L., O. Manor, and C. Power. 2004. "Early Environment and Child-to-Adult Growth Trajectories in the 1958 British Birth Cohort." *American Journal of Clinical Nutrition* 80:185–92.
- Lynn, R. 1989. "A Nutrition Theory of the Secular Increases in Intelligence, Positive Correlation Between Height, Head Size and IQ." *British Journal of Educational Psychology* 59:372–77.
- Magnusson, P.K.E., F. Rasmussen, and U.B. Gyllenstein. 2006. "Height at Age 18 Years Is a Strong Predictor of Attained Education Later in Life: Cohort Study of Over 950,000 Swedish Men." *International Journal of Epidemiology* 35:658–63.
- Maluccio, J.A., J. Hoddinott, J.R. Behrman, R. Martorell, A.R. Quisumbing, and A.D. Stein. 2009. "The Impact of Improving Nutrition During Early Childhood on Education Among Guatemalan Adults." *Economic Journal* 119:734–63.
- Marmot, M., M. Shipley, E. Brunner, and H. Hemingway. 2001. "Relative Contribution of Early Life and Adult Socioeconomic Factors to Adult Morbidity in the Whitehall II Study." *Journal of Epidemiology and Community Health* 55:301–307.
- McMillen, I.C. and J.S. Robinson. 2005. "Developmental Origins of the Metabolic Syndrome: Prediction, Plasticity, and Programming." *Physiological Reviews* 85:571–633.
- Nilsson, C., M. Niklasson, E. Eriksson, P. Bjorntorp, and A. Holmang. 1998. "Imprinting of Female Offspring With Testosterone Results in Insulin Resistance and Changes in Body Fat Distribution at Adult Age in Rats." *Journal of Clinical Investigation* 101:74–78.
- Ozanne, S.E. 2001. "Metabolic Programming in Animals." *British Medical Bulletin* 60:143–52.
- Persico, N., A. Postlewaite, and D. Silverman. 2004. "The Effect of Adolescent Experience on Labor Market Outcomes: The Case of Height." *Journal of Political Economy* 112:1019–53.
- Phillips, D.I. 2001. "Fetal Growth and Programming of the Hypothalamic-Pituitary-Adrenal Axis." *Clinical and Experimental Pharmacology and Physiology* 28:967–70.
- Räikkönen, K., T. Forsén, M. Henriksson, E. Kajantie, K. Heinonen, A. Pesonen, J.T. Leskinen, I. Laaksonen, C. Osmond, D.J.P. Barker, and J.G. Eriksson. 2009. "Growth Trajectories and Intellectual Abilities in Young Adulthood, The Helsinki Birth Cohort Study." *American Journal of Epidemiology* 170:447–55.
- Rutter, M. and the English and Romanian Adoptees (ERA) Study Team. 1998. "Developmental Catch-up and Deficit Following Adoption After Severe Global Early Privation." *Journal of Child Psychology and Psychiatry* 39:465–76.
- Saenger, P., P. Czernichow, I. Hughes, and E.O. Reiter. 2007. "Small for Gestational Age: Short Stature and Beyond." *Endocrine Reviews* 28:219–51.
- Samuelsson, A.M., I. Ohrn, J. Dahlgren, E. Eriksson, B. Angelin, B. Folkow, and A. Holmang. 2004. "Prenatal Exposure to Interleukin-6 Results in Hypertension and Increased Hypothalamic-Pituitary-Adrenal Axis Activity in Adult Rats." *Endocrinology* 145:4897–911.
- Smith, J.P. 2009. "The Impact of Childhood Health on Adult Labor Market Outcomes." *The Review of Economics and Statistics* 91:478–89.

- Steckel, R.H. 1995. "Stature and the Standard of Living." *Journal of Economic Literature* 33: 1903–40.
- Sundet, J.M., K. Tambs, J.R. Harris, P. Magnus, and T.M. Torjussen. 2005. "Resolving the Genetic and Environmental Sources of the Correlation Between Height and Intelligence: A Study of Nearly 2600 Norwegian Male Twin Pairs." *Twin Research and Human Genetics* 8:307–11.
- Walker, S.P., S.M. Chang, C.A. Powell, and S.M. Grantham-McGregor. 2005. "Effects of Early Childhood Psychosocial Stimulation and Nutritional Supplementation on Cognition and Education in Growth-Stunted Jamaican Children: Prospective Cohort Study." *Lancet* 366:1804–807.