Answers to Hour Examination #2, Chemistry 302X-302A, 2005

1.

- (a) Aldol followed by reductions:
- 1. H₃O⁺/H₂O (aldol) 2. H₂/Pd/C (reduce) 3. NH₂NH₂/HOCH₂CH₂OH/heat (Wolff-Kischner you could also use the Clemmenson)

or

- 1. LiAlH₄ 2. H₂O 3. HBr 4. Li (makes cyclohexyllithium) 5. react with cyclohexanone 6. H₂O/H₃O⁺ (protonate and eliminate). 7. H₂/Pd/C
- (b) Aldol followed by Michael:
- 1. H₃O⁺/H₂O 2. KCN, H₂O
- (c) Knoevenagel:
- 1. (CH3OOC)2CH2/CH3O- Na+
- 2. neutralize

2.

- (a) 1. LDA 2. CH₃I (an enamine process is also fine)
- (b) 1. CH₃Li 2. H₂O 3. H₃O⁺/H₂O (E1) 4. BH₃ 5. HOOH/HO⁻ 6. KMnO₄

3.

4. It is clear where the new bond joining the two pieces must be formed.

$$CH_3$$
 CH_3

But it is equally clear, we hope, that there is no acidic hydrogen available for removal.

The role of the amine is to Michael in, thus generating the necessary resonancestabilized intermediate.

$$CH_3$$
 CH_3
 CH_3
 CH_3

this intermediate is resonance stabilized

Now the junction bond can be formed.

5. No magic methyl shifts here - this must be an open-close problem. Let's do it backward: as we know that all - -unsaturated carbonyl compounds come from aldols, just deconstruct the product to the open compound we must make.

here are the steps sketched out for the closure of the dialdehyde to the product - not every keto-enol eq. is shown in detail here:

Here is the formation of the dialdehyde from starting material:

Michael.