

D.C. Economists Minicourse –
What's New in Econometrics: Time Series

Lecture 10

November 14, 2008

Forecast Assessment

Outline

1. Why Forecast?
2. Forecasting basics
3. Estimating Parameters for forecasting
4. Forecast Assessment
 - (a) Comparing Forecasts/Forecasters
 - (b) Comparing Forecasts/Models

1. Why Forecast?

- (a) You want to know about the future (SPF, Financial Markets)
- (b) You want to evaluate a model

Why use forecasting methods to evaluate models?

- (i) Less prone to data mining, in-sample overfitting (indeed, in-sample overfitting leads to out-of-sample “underfitting”)
- (ii) Instability
- (iii) In-sample methods too difficult

Recently developed forecast assessment methods have focused on both of these (and it is useful to keep goals in mind as we discuss methods)

2. Forecasting Basics

Y_{t+h} : variable to be forecast

X_t : vector of variables used to make forecast (typically would include current and lags of Y_t and other variables).

$f_{t+h/t}$: Forecast of Y_{t+h} made a time t .

$e_{t+h} : Y_{t+h} - f_{t+h/t}$

$L(e)$: Loss associated with the error

“Risk” associated with f : $E(L(e))$.

(a) Minimum MSE forecasts: Loss quadratic

$L(e) = a + be^2$ and Risk is Mean Squared Error (MSE). Goal is to find minimum MSE (MMSE) forecast.

Key Result: MMSE forecast is the regression function: $f_{t+h/t} = E(Y_{t+h}|X_t)$

Some properties of MMSE forecasts:

(1) $E(e_{t+h} | X_t) = 0$, so that $E(e_{t+h}X_t) = 0$

(2) If X_t includes current and past values of Y_t , then it implicitly includes current and lagged values of e_t . Thus $E(e_{t+h}e_t) = 0$, so that e_{t+h} follows an MA($h-1$) process.

(3) $Y_{t+h} = f_{t+h/t} + e_{t+h}$, where the two rhs terms are uncorrelated. Thus, $\sigma_Y^2 = \sigma_f^2 + \sigma_e^2$, and $\sigma_Y^2 \geq \sigma_f^2$.

(b) More than one forecast:

Let f^1 and f^2 denote two forecasts of Y . Let $f^3 = \beta_0 + \beta_1 f^1 + \beta_2 f^2$ denote a third forecast.

Question: How should f^1 and f^2 be “combined” to form f^3 (what values of β should be chosen)?

Answer: MMSE forecasts are regressions, so β 's are given by the (population) values from the linear regression of Y_{t+h} onto $f_{t+h/t}^1$ and $f_{t+h/t}^2$. (Bates and Granger (1969), Granger and Ramanathan (1984)).

The extension to $n \geq 2$ forecasts is obvious.

References: see Timmermann (2006)

Problem: As a practical matter you must use estimates of the β 's from a sample regression.

Forecast combining “puzzle”: When the number of forecasts to be combined (n) is even moderately large, forecasts constructed using estimated β 's don't perform very well. Better to use ad hoc averages like sample means, medians, trimmed means, “consensus” forecasts and so forth. (Large literature surveyed in Timmermann.) JHS will say more about this in Lecture 12.

(c) Other Loss Functions:

Granger (1969) and Christoffersen and Diebold (1997).

If $\text{Risk}(f) = E(L(Y-f))$, and $Y_{t+h} | X_t \sim N(\mu_{t+h/t}, \sigma_{t+h/t}^2)$, then the optimal forecast is $f_{t+h/t} = \mu_{t+h/t} + \alpha(\sigma_{t+h/t}^2)$.

Proof: Write $Y_{t+h} = \mu_{t+h/t} + e_{t+h}$ and $f_{t+h/t} = \mu_{t+h/t} + \alpha_{t+h/t}$. Then

$E(L(Y-f)) = E(L(e - \alpha))$, and the probability density of $L(e - \alpha)$ depends only on $\sigma_{t+h/t}^2$.

Implication: In condition homoskedastic Gaussian world, optimal forecasts are MMSE forecasts $(\mu_{t+h/t}) + \text{constant}$.

Elliott, Komunjer and Timmermann (2005):

$$L(Y-f) = [\alpha + (1-2\alpha)\mathbf{1}(Y-f < 0)] |Y-f|^p$$

Non-quadratic, non-symmetric, ...

(They study properties of optimal linear forecasters $f_{t+h/t} = \theta'X_t$, with an aim to robustifying tests for forecast efficiency.)

Estimating Parameters for Use in Forecasting Models:

(1) Should you use MLE or other estimators?

Example: AR(1) model: $y_t = \phi y_{t-1} + \varepsilon_t$

Goal: forecast y_{t+2}

Optimal forecast: $f_{t+2/t} = \beta y_t$, where $\beta = \phi^2$

Two ways to forecast y_{t+2}

“*Iterated*”: Estimate ϕ from $y_t = \phi y_{t-1} + \varepsilon_t$, and use $\hat{f}_{t+2/t}^{iterated} = \hat{\phi}^2 y_t$

“*Direct*”: Estimate β from $y_t = \beta y_{t-2} + u_t$, and use $\hat{f}_{t+2/t}^{direct} = \hat{\beta} y_t$

Pros and Cons:

$\hat{\phi}$ is MLE, and is “efficient” under AR(1), but what if model is misspecified?

$\hat{\beta}$ has larger variance than $\hat{\phi}$ under correct specification, but is robust to misspecification (for the class of forecasts under consideration).

Literature: Cox (1961) to Schorfheide (2005) (survey: Bhansali (1999))

Empirical Comparison: Marcelino, Stock and Watson (2006)

170 Monthly U.S. Macro Series, 1959-2002

Pseudo-out-of-sample forecasts (POOS)

AR and Bivariate VAR for $h = 3, 6, 12, 24$

$$\text{Relative pseudo-out-of-sample MSE} = \frac{\sum (e_{t+h}^{\text{direct}})^2}{\sum (e_{t+h}^{\text{iterated}})^2}$$

Lag Length	Horizon			
	3	6	12	24
AR(4)	0.99	0.99	1.00	1.05
AR(12)	1.01	1.01	1.03	1.10
AR(BIC)	0.98	0.97	0.99	1.05
AR(AIC)	1.00	1.01	1.02	1.09

Across Lag-Length Methods: AIC Iterated seemed to work best.

Estimating Parameters for Use in Forecasting Models:

(2) Should you use “Real-Time Data”?

$$y_{t+h} = \beta'x_t + u_{t+h}$$

Data Revisions in y and x . Issues to think about:

(a) y : what the goal of forecasting (first release, final release, ...)

(b) x : real-time forecasting using real-time data, so it seems that x^{initial} should be used in regressions to estimate β .

Key question: is projection of y on x the same as projection of y on x^{initial} ?

$x = x^{\text{initial}} + x^{\text{revision}}$... is revision “News” or “Noise” in the Mankiw, Runkle, and Shapiro (1984) dichotomy.

Forecast Assessment – Part 1 (Evaluating Forecasts/Forecasters ... Not Models)

(a) Mincer-Zarnowitz (1969) Regressions: $Y_{t+h} = \alpha + \beta f_{t+h/t} + \gamma W_t + u_{t+h}$

If $f_{t+h/t}$ is MMSE forecast then $\alpha = 0$, $\beta = 1$, $\gamma = 0$. Test all or subset.

Inference Issues:

(i) $h > 1$, $u_{t+h} \sim \text{MA}(h-1)$ under H_0 . Use HAC SEs.

(ii) If Y_t is persistent (say $I(1)$), then so is f , so “unit-root” regression inference problems. Easy fix: $(Y_{t+h} - Y_t) = \alpha + \beta(f_{t+h/t} - Y_t) + \gamma W_t + u_{t+h}$.

(iii) h large, u is very persistent. HAC works poorly. (Richardson and Stock (1989)).

(b) Combining or “Encompassing” Regressions

2 forecasts f^1 and f^2 .

Forecast combining regression: $Y_{t+h} = \beta_0 + \beta_1 f_{t+h/t}^1 + \beta_2 f_{t+h/t}^2 + u_{t+h}$

If f^1 is MMSE forecast, then $\beta_0 = \beta_2 = 0$ and $\beta_1 = 1$.

Inference Issues: – Same as MZ regressions. Also, you might want to make alternative more parsimonious imposing $\beta_0 = 0$ and $\beta_1 + \beta_2 = 1$.

Imposing this constraint yields

$$Y_{t+h} - f_{t+h/t}^1 = \beta_2(f_{t+h/t}^2 - f_{t+h/t}^1) + u_{t+h}.$$

(c) Loss-Function Tests

2 forecasts f^1 and f^2 with forecast errors e^1 and e^2 .

Question: Is $E(L(e^1)) = E(L(e^2))$ (or $<$ or $>$)?

Testing using quadratic loss. Realized forecast errors e_t^1 and e_t^2 .

Question: $E[(e_t^1)^2] = E[(e_t^2)^2]$ or $E[(e_t^1)^2 - (e_t^2)^2] = 0$?

Sample moments of differences: $\bar{d} = \frac{1}{T} \sum_{t=1}^T d_t$, where $d_t = \{(e_t^1)^2 - (e_t^2)^2\}$

Is \bar{d} statistically significantly different from zero.

Inference Issues: (i) $h > 1$, serial correlation. Use HAC se's.

Refs: Diebold and Mariano (1995), see West (2006) for history

(d) Loss function tests with many competing forecasts:

f^1 is a benchmark model (say, random walk)

$f^k, k = 1, \dots, n$ are n competing models

Question: Do any of the n competing models dominate the benchmark?

$\bar{d}_k = \frac{1}{T} \sum_{t=1}^T d_t^k$, where $d_t^k = \{(e_t^1)^2 - (e_t^k)^2\}$, so that \bar{d}_k is the sample MSE

improvement for model k over benchmark prediction.


Test Statistic: $RC = \max_k \bar{d}_k$ (RC for “Reality Check”)

White (2000) derives limiting distribution of RC under null that benchmark is optimal, so that critical values can be computed.

(See Hansen (2005) for a refinement.)

Density Forecasts:

Chart 2 CPI inflation projection based on market interest rate expectations



Evaluating Density Forecasts

Ref: Diebold, Gunther, Tay (1998)

Key Insight: Suppose Y has CDF F . Then $U = F(Y) \sim U[0,1]$. (Recall random number generators often use $Y = F^{-1}(U)$.)

Thus, if $F_{t+h/t}$ is the conditional CDF of $Y_{t+h/t}$ then $U_{t+h} = F_{t+h/t}(Y_{t+h})$ should be $U[0,1]$ and U_{t+h} should be independent of data dated t and earlier.

Interval Forecasts: “Provide a 95% ‘confidence interval’” for GDP growth in the final quarter of 2008. (ref: Christoffersen (1998)).

Forecast Assessment – Part 2 (Evaluating Models using POOS)

Key References: West (1996, 2006)

Key Statistical Difference Between this and Part 1: Explicitly account for sampling variability in estimated model parameters.

Setup: Model 1 has forecasts $f^1(\theta_1)$ and Model 2 has forecasts $f^2(\theta_2)$. The forecasts that are evaluated are based on estimated models $f^1(\hat{\theta}_1)$ and $f^2(\hat{\theta}_2)$.

Pseudo-Out-Of Sampling (POOS) Forecasting Strategy:

Sample Size $T = R + P$. Last P periods used for ‘Prediction’ (construction of pseudo-out-of-sample forecasts).

Strategy:

- (i) Estimate θ using observations 1: R , $\hat{\theta}_R$
- (ii) Forecast Y_{R+h} using data through R and $\hat{\theta}_R$.

then (recursive POOS)

- (iii) Estimate θ using observations 1: $R+1$, $\hat{\theta}_{R+1}$
- (iv) Forecast Y_{R+1+h} using data through $R+1$ and $\hat{\theta}_{R+1}$.

or (rolling POOS)

- (iii) Estimate θ using observations 2: $R+1$, $\hat{\theta}_{R+1}$
- (iv) Forecast Y_{R+1+h} using data through $R+1$ and $\hat{\theta}_{R+1}$.

...

Note (and this is important):

The Question of interest is:

$$(i) E[L(Y - f^1(\theta_1))] = E[L(Y - f^2(\theta_2))] \text{ (or } < \text{ or } > \text{)}$$

NOT:

$$(ii) E[L(Y - f^1(\hat{\theta}_1))] = E[L(Y - f^2(\hat{\theta}_2))] \text{ (or } < \text{ or } > \text{) (but I will return to this)}$$

Many interesting cases in which

$$E[L(Y - f^1(\hat{\theta}_1))] < E[L(Y - f^2(\hat{\theta}_2))], \text{ but } E[L(Y - f^1(\theta_1))] > E[L(Y - f^2(\theta_2))]$$

(Example: Random Walk in Exchange Rates ... Engel and West (2005), Clark and West (2006), Rossi (2005).)

Focus here is on risk comparisons. Related issues arise in combining tests.

A Complication: Nested vs. Non-nested models

Nested Models:

Model 1: $y_{t+1} = x_t' \beta + \varepsilon_{t+1}$ is a special case of

Model 2: $y_{t+1} = x_t' \beta + z_t' \gamma + e_{t+1}$

Non-Nested Models

Model 1: $y_{t+1} = x_t' \beta + \varepsilon_{t+1}$ is not a special case of

Model 2: $y_{t+1} = z_t' \gamma + e_{t+1}$

Non-Nested Models (West (1996)):

Model 1: $y_{t+1} = x_t\beta + \varepsilon_{t+1}$ (scalar x , $h = 1$ -step-ahead for convenience)

True Forecast: $x_t\beta$

True Forecast error: ε_{t+1}

Estimated Forecast: $x_t\hat{\beta}_t$

Estimated error: $Y_{t+1} - x_t\hat{\beta}_t = \hat{\varepsilon}_{t+1} = \varepsilon_{t+1} + x_t(\hat{\beta}_t - \beta)$

Model 2: $y_{t+1} = z_t\gamma + e_{t+1}$

True Forecast: $z_t\gamma$

True Forecast error: e_{t+1}

Estimated Forecast: $z_t\hat{\gamma}_t$

Estimated error: $Y_{t+1} - z_t\hat{\gamma}_t = \hat{e}_{t+1} = e_{t+1} + z_t(\hat{\gamma}_t - \gamma)$

Recall that Loss Function Test looks at averages of $d_t = (\varepsilon_t^2 - e_t^2)$ over the prediction period. Here, we must use $\hat{d}_t = (\hat{\varepsilon}_t^2 - \hat{e}_t^2)$. How are the sample averages of \hat{d}_t and d_t related?

A Calculation:

$$\begin{aligned}
 \frac{1}{\sqrt{P}} \sum_{t=R+1}^T (\hat{\varepsilon}_t^2 - \hat{e}_t^2) &= \frac{1}{\sqrt{P}} \sum_{t=R+1}^T (\varepsilon_t^2 - e_t^2) \\
 &+ \frac{1}{\sqrt{P}} \sum_{t=R+1}^T (\hat{\beta}_{t-1} - \beta)^2 x_{t-1}^2 + 2 \frac{1}{\sqrt{P}} \sum_{t=R+1}^T (\hat{\beta}_{t-1} - \beta) x_{t-1} \varepsilon_t \\
 &+ \frac{1}{\sqrt{P}} \sum_{t=R+1}^T (\hat{\gamma}_{t-1} - \gamma)^2 z_{t-1}^2 + 2 \frac{1}{\sqrt{P}} \sum_{t=R+1}^T (\hat{\gamma}_{t-1} - \gamma) z_{t-1} e_t \\
 &= \frac{1}{\sqrt{P}} \sum_{t=R+1}^T (\varepsilon_t^2 - e_t^2) + o_p(1)
 \end{aligned}$$

where (from West (1996)) the final inequality holds when the $E(\varepsilon_t x_{t-1}) = E(e_t z_{t-1}) = 0$ (and some technical assumptions are satisfied).

Bottom Line: $\frac{1}{\sqrt{P}} \sum_{t=R+1}^T (\hat{\varepsilon}_t^2 - \hat{e}_t^2) \approx \frac{1}{\sqrt{P}} \sum_{t=R+1}^T (\varepsilon_t^2 - e_t^2)$ (so sampling error if $\hat{\beta}$ and $\hat{\gamma}$ do not matter.

Nested Models: McCracken (2000), Clark and McCracken (2001)

Identity (from last page):

$$\begin{aligned} \sum_{t=R+1}^T (\hat{\varepsilon}_t^2 - \hat{e}_t^2) &= \sum_{t=R+1}^T (\varepsilon_t^2 - e_t^2) \\ &+ \sum_{t=R+1}^T (\hat{\beta}_{t-1} - \beta)^2 x_{t-1}^2 + 2 \sum_{t=R+1}^T (\hat{\beta}_{t-1} - \beta) x_{t-1} \varepsilon_t \\ &+ \sum_{t=R+1}^T (\hat{\gamma}_{t-1} - \gamma)^2 z_{t-1}^2 + 2 \sum_{t=R+1}^T (\hat{\gamma}_{t-1} - \gamma) z_{t-1} e_t \end{aligned}$$

Non-Nested Model: $\sum_{t=R+1}^T (\varepsilon_t^2 - e_t^2) \sim O_p(P^{1/2})$ dominates rhs

Nested Models: $y_{t+1} = x_t' \beta + \varepsilon_{t+1}$ and $y_{t+1} = x_t' \beta + z_t' \gamma + e_{t+1}$

under equal loss, models are the same, so that $\varepsilon_t = e_t$. First term vanishes.

$$\begin{aligned}
\sum_{t=R+1}^T (\hat{\varepsilon}_t^2 - \hat{e}_t^2) &= 0 \\
&+ \sum_{t=R+1}^T (\hat{\beta}_{t-1} - \beta)^2 x_{t-1}^2 + 2 \sum_{t=R+1}^T (\hat{\beta}_{t-1} - \beta) x_{t-1} \varepsilon_t \\
&+ \sum_{t=R+1}^T (\hat{\gamma}_{t-1} - \gamma)^2 z_{t-1}^2 + 2 \sum_{t=R+1}^T (\hat{\gamma}_{t-1} - \gamma) z_{t-1} e_t
\end{aligned}$$

This is harder ... Papers by McCracken and Clark and McCracken attack this by studying behavior of terms on rhs.

Key results: Limits are (somewhat complicated/messy) functions of normals.

Implication: Can use parametric bootstrap (Gaussian errors) to approximate limiting distribution, compute critical values and so forth.

An Important Special Case: f^1 is a random walk forecast, f^2 is nested within the random walk. (Clark and West (2006)).

Let y_t denote the first difference of a variable of interest (e.g., an exchange rate). Under the null, y_t is mds, while under the alternative it can be predicted by x_t .

Model 1: $y_{t+1} = \varepsilon_{t+1}$

Forecast 1: $\hat{f}_{t+1/t}^1 = 0$

Errors (under H_0): $\hat{e}_{t+1}^1 = \varepsilon_{t+1}$

Model 2: $y_{t+1} = x_t' \beta + \varepsilon_{t+1}$

Forecast 2: $\hat{f}_{t+1/t}^2 = x_t' \hat{\beta}_t$

Errors (under H_0): $\hat{e}_{t+1}^2 = \varepsilon_{t+1} - x_t' \hat{\beta}_t$

MSPE (under H_0):

$$\frac{1}{P} \sum_{t=R+1}^P \varepsilon_{t+1}^2$$

$$\frac{1}{P} \sum_{t=R+1}^P \varepsilon_{t+1}^2 + \frac{1}{P} \sum_{t=r+1}^P (x_t' \hat{\beta}_t)^2 - 2 \frac{1}{P} \sum_{t=r+1}^P \varepsilon_{t+1} x_t' \hat{\beta}_t$$

Expectation:

$$\sigma_\varepsilon^2$$

$$\sigma_\varepsilon^2 + E \left(\frac{1}{P} \sum_{t=r+1}^P (x_t' \hat{\beta}_t)^2 \right)$$

Thus, under RW null

$$E(\text{MSE for RW}) = E(\text{MSE for alternative}) - E\left(\frac{1}{P} \sum_{t=r+1}^P (x_t' \hat{\beta}_t)^2\right)$$

RW does not suffer from overfitting ($x_t' \hat{\beta}_t$) and should forecast better than alternative ... and ... the “overfitting” term can be estimated:

$$E\left(\frac{1}{P} \sum_{t=r+1}^P (x_t' \hat{\beta}_t)^2\right) \approx \frac{1}{P} \sum_{t=r+1}^P (x_t' \hat{\beta}_t)^2 = \frac{1}{P} \sum_{t=r+1}^P (f_{t/t-1}^2)^2$$

Clark-West test: standardized version of $\hat{\sigma}_1^2 - \left(\hat{\sigma}_2^2 - \frac{1}{P} \sum_{t=r+1}^P (f_{t/t-1}^2)^2\right)$,

where $\hat{\sigma}_1^2$ and $\hat{\sigma}_2^2$ are the POOS mean square prediction errors for f^1 (the RW) and f^2 .

What is the distribution of this difference?

$$\sqrt{P} \left(\hat{\sigma}_1^2 - \left(\hat{\sigma}_2^2 - \frac{1}{P} \sum_{t=r+1}^P (f_{t/t-1}^2)^2 \right) \right) = \frac{1}{\sqrt{P}} \sum_{t=r+1}^P \varepsilon_t f_{t/t-1}^2$$

And $\varepsilon_t f_{t/t-1}^2$ is a mds under the null. Maybe a normal limit?

Caution: (example) $x_t = 1$, $f_{t/t-1}^2 = \hat{\beta}_{t-1} = \frac{1}{t-1} \sum_{i=1}^{t-1} y_i \stackrel{H_0}{=} \frac{1}{t-1} \sum_{i=1}^{t-1} \varepsilon_i$, and

$\varepsilon_t f_{t/t-1}^2 \stackrel{H_0}{=} \frac{1}{t-1} \varepsilon_t \sum_{i=1}^{t-1} \varepsilon_i$. Remember unit root AR Model, numerator of $\hat{\rho} - \rho$ is

$$\sum_{t=1}^T \varepsilon_t \sum_{i=1}^{t-1} \varepsilon_i.$$

CW: Using “Rolling” estimate of β based on R observations to limit this dependence. (Where R is fixed and not too large – Giacomini and White (2001), discussed below.)

CW Empirical Example:

Exchange Rates (\$'s) Monthly: Canada, Japan, Switzerland, UK

POOS Period 1990-2003 ($P = 166$), use $R = 120$.

$x = (1, \text{1-month interest differential})$

Table 6
Forecasts of monthly changes in US Dollar exchange rates

(1) Country	(2) Prediction sample	(3) $\hat{\sigma}_1^2$	(4) $\hat{\sigma}_2^2$	(5) adj.	(6) $\hat{\sigma}_2^2$ -adj.	(7) MSPE-adjusted $\hat{\sigma}_1^2 - (\hat{\sigma}_2^2 - \text{adj.})$	(8) MSPE-normal $\hat{\sigma}_1^2 - \hat{\sigma}_2^2$	(9) CCS
Canada	1990:1– 2003:10	2.36	2.32	0.09	2.22	0.13 (0.08) <i>1.78**</i>	0.04 <i>0.54††</i>	<i>3.67</i>
Japan	1990:1– 2003:10	11.32	11.55	0.75	10.80	0.53 (0.43) <i>1.24</i>	−0.23 <i>−0.52</i>	<i>5.23*</i>
Switzerland	1985:1– 2003:10	12.27	12.33	0.96	11.37	0.90 (0.48) <i>1.88**</i>	−0.06 <i>−0.13</i>	<i>2.43</i>
U.K.	1985:1– 2003:10	9.73	10.16	0.44	9.72	0.01 (0.33) <i>0.03</i>	−0.43 <i>−1.27</i>	<i>0.78</i>

1. Forecast Assessment – Part 1 (Evaluating Forecasts/Forecasters)

$$E[L(Y - f^1)] = E[L(Y - f^2)] \text{ (or } < \text{ or } > \text{)}$$

2. Forecast Assessment – Part 2 (Evaluating Models using POOS forecasting)

$$E[L(Y - f^1(\theta_1))] = E[L(Y - f^2(\theta_2))] \text{ (or } < \text{ or } > \text{)}$$

3. Forecasting Assessment – Part 3 (Evaluating Forecasting Models using POOS forecasting)

$$E[L(Y - f^1(\hat{\theta}_1))] = E[L(Y - f^2(\hat{\theta}_2))] \text{ (or } < \text{ or } > \text{)}$$

Giacomini and White (2006)

Giacomini and White (2006):

	Data	Forecasting Procedure	Forecast
Forecasting Model 1:	$\{x_i, y_i\}_{i=1}^t$	$\hat{\theta}_1$ and whatever	$f_{t+1/t}^1$
Forecasting Model 2:	$\{x_i, y_i\}_{i=1}^t$	$\hat{\theta}_2$ and whatever	$f_{t+1/t}^2$

Question: $E\left[L(f_{t+1/t}^1) - L(f_{t+1/t}^2) \mid g_t\right] = 0$ (or $<$ or $>$)

(Using time t information, can I predict which forecast will have smaller loss?)

GMM Problem ... $E\left[\left(L(f_{t+1/t}^1) - L(f_{t+1/t}^2)\right)q(g_t)\right] = 0$

$$\text{Moment: } E \left[\left(L(f_{t+1/t}^1) - L(f_{t+1/t}^2) \right) q(g_t) \right] = 0$$

Forecasting Model 1: $\{x_i, y_i\}_{i=1}^t \rightarrow \boxed{\hat{\theta}_1 \text{ and whatever}} \rightarrow f_{t+1/t}^1$

Forecasting Model 2: $\{x_i, y_i\}_{i=1}^t \rightarrow \boxed{\hat{\theta}_2 \text{ and whatever}} \rightarrow f_{t+1/t}^2$

Inference Issues:

(1) Stationarity: using rolling samples $\{x_i, y_i\}_{i=t-R}^t$

(2) h -step-ahead prediction: HAC (allowing for h dependence)

(3) Unconditional comparisons ($q(g_t) = 1$): Much more dependence (at least R) ... method is (arguably) less useful. (Note: This is a different question than the question that motivated their analysis.)