

Using Imprecise Measures to Study Component and System Reliability

Kimberly F. Sellers
Department of Statistics
Carnegie Mellon University

January 17, 2003

Outline

- Binary coherent systems with precise classification
 - Definition
 - Reliability
- Binary state system with imprecise classification
 - Membership functions and probability of fuzzy sets
 - Reliability of degraded components and systems

Binary Coherent Systems

- Range of states, $S=\{0, 1\}$
- State of the i th component,

$$X_i = \begin{cases} 1, & \text{if component } i \text{ functions} \\ 0, & \text{if component } i \text{ fails} \end{cases}$$

- Structure function,

$$\phi = \phi(\mathbf{X}) = \begin{cases} 1, & \text{if the system functions} \\ 0, & \text{if the system fails.} \end{cases}$$

Binary Coherent Systems (cont.)

- ϕ is a *binary coherent system* if
 1. ϕ is nondecreasing
 2. each component is relevant, i.e. $\forall i, \exists(\cdot_i, \mathbf{X}) \ni$

$$\phi(1_i, \mathbf{X}) = 1 \text{ and } \phi(0_i, \mathbf{X}) = 0.$$

Binary Coherent Systems (cont.)

Basic examples of binary coherent systems:

- Series:

$$\phi(\mathbf{X}) = \prod_{i=1}^n X_i$$

- Parallel (not stand-by):

$$\phi(\mathbf{X}) = \prod_{i=1}^n X_i = 1 - \prod_{i=1}^n (1 - X_i)$$

- k -out-of- n :

$$\phi(\mathbf{X}) = \begin{cases} 1, & \text{if } \sum_{i=1}^n X_i \geq k \\ 0, & \text{if } \sum_{i=1}^n X_i < k \end{cases}$$

Reliability of Binary Coherent Systems

- $X_i \mid p_i \sim \text{Bernoulli}(p_i)$
 - $p_i = P(X_i = 1 \mid p_i)$ – reliability of component i
 - $h(\mathbf{p}) = P(\phi(\mathbf{X}) = 1 \mid \mathbf{p})$ – reliability of ϕ
 - System reliability is determined by the reliability of its components
- Assumptions:
 - Given \mathbf{p} , $X_i \perp X_j \forall i \neq j$.
 - Given p_i , $X_i \perp p_j \forall j \neq i$.

Reliability of Binary Coherent Systems (cont.)

- Series system:

$$h_S(\mathbf{p}) = P(\phi(\mathbf{X}) = 1 \mid \mathbf{p}) = \prod_{i=1}^n p_i$$

- Parallel (not stand-by) system:

$$h_P(\mathbf{p}) = \prod_{i=1}^n p_i = 1 - \prod_{i=1}^n (1 - p_i)$$

- k -out-of- n system:

$$h_K(\tilde{p}) = \sum_{j=k}^n \binom{n}{j} \tilde{p}^j (1 - \tilde{p})^{n-j}$$

if all components are identical with reliability $p_i \equiv \tilde{p}$.

Binary State Systems with Imprecise Classification

- Membership functions and probability of fuzzy sets
- Reliability of degraded components and systems

Membership Function, $\mu_A(x)$

- For each x , $0 \leq \mu_A(x) \leq 1$ describes a belief of containment (membership) in a category A .
- If $\mu_A(x) = 0$ or 1 , then A is a precise (i.e., crisp) set. Otherwise, A is a fuzzy set.

Ex. 1: $A_1 = \{x \in (1, 2, \dots, 10) \mid x \geq 7\}$

x	1	2	3	4	5	6	7	8	9	10
$\mu_{A_1}(x)$	0	0	0	0	0	0	1	1	1	1

Ex. 2: $A_2 = \{x \in (1, 2, \dots, 10) \mid x \text{ is large}\}$

x	1	2	3	4	5	6	7	8	9	10
$\mu_{A_2}(x)$	0	0	0	0	0	0.2	0.5	0.9	1	1

Membership Function, $\mu_A(x)$

- Operations:
 1. $\mu_{A \cup B}(x) = \max(\mu_A(x), \mu_B(x))$
 2. $\mu_{A \cap B}(x) = \min(\mu_A(x), \mu_B(x))$
 3. $\mu_{A^c}(x) = 1 - \mu_A(x)$
- Definitions parallel Lukasiewicz's (1930) definitions for the conjunction, disjunction and complement of many-valued propositions
- Bellman and Giertz (1973) showed that Operations 1 and 2 are unique with respect to constructing an algebra of sets; Operation 3 is not unique but "appears reasonable".

Probability of Fuzzy Sets

Due to Bement et. al. (2000)

- Based on two premises: $\mu_{\tilde{A}}(x)$ is interpreted as a likelihood function, and fuzzy sets arise from boundary uncertainty
- $\pi_D(x \in \tilde{A})$ is D 's probability that $x \in \tilde{A}$ if it is known that $X = x$
- $\mu_{\tilde{A}}(x)$ is additional information (expert testimony) provided by expert, Z

$$P_D(X \in \tilde{A}; \mu_{\tilde{A}}(x)) \propto \int_x \mu_{\tilde{A}}(x) \pi_D(x \in \tilde{A}) dP_D(x) \quad (1)$$

Probability of Fuzzy Sets (cont.)

- D needs to specify two probability measures, P_D and π_D
 - $(\Omega, \mathcal{F}, P_D)$, where Ω is the set of all possible outcomes of the random phenomenon under study
 - $(\Omega, \mathcal{F}, \pi_D)$, where Ω consists of two outcomes $x \in \tilde{A}$ or $x \notin \tilde{A}$
 - $\pi_D(x \in \tilde{A}) = 1 - \pi_D(x \notin \tilde{A})$

- To evaluate the constant of proportionality, D needs to evaluate

$$P_D(X \notin \tilde{A}; \mu_{\tilde{A}}(x)) \propto \int_x \mathcal{L}_D(X \notin \tilde{A}; \mu_{\tilde{A}}(x)) \pi_D(x \notin \tilde{A}) dP_D(x) \quad (2)$$

- $\mathcal{L}_D(X \notin \tilde{A}; \mu_{\tilde{A}}(x))$ not necessarily $1 - \mu_{\tilde{A}}(x)$

Components in Vague Binary States


- Let $X =$ the state of a component at time $\tau > 0$
- X takes values in $\mathcal{S} = \{x; 0 \leq x \leq 1\}$
- Consider $\mathcal{S}_1 \subset \mathcal{S}$, where $\mathcal{S}_1 = \{x; x \text{ is a "desirable" state}\}$ and $\mu_1(x)$ membership function
- If $\mathcal{S}_2 \subset \mathcal{S}$ was defined as

$$\mathcal{S}_2 = \{x; x \text{ is an "undesirable" state}\},$$


then \mathcal{S}_1^C need not be \mathcal{S}_2 unless $\mu_2(x) = 1 - \mu_1(x)$, where $\mu_2(x)$ is the membership function of \mathcal{S}_2

Components in Vague Binary States (cont.)

- $\mu_2(x)$ need not be symmetrical to $\mu_1(x)$



Symmetric case



Asymmetric case

- In general, \mathcal{S}_1^C need not be \mathcal{S}_2 and vice versa, unless \mathcal{S}_1 and \mathcal{S}_2 are precise sets.

Components in Vague Binary States (cont.)

- For $\mathcal{S}_1 = \{x; x \text{ is a "desirable" state}\}$ and $\mu_1(x)$ specified, we define component reliability as

$$P_D(X \in \mathcal{S}_1; \mu_1(x)) \propto \int_x \mu_1(x) \pi_D(x \in \mathcal{S}_1) dP_D(x). \quad (3)$$

- With $\mathcal{S}_2 = \{x; x \text{ is an "undesirable" state}\}$, and $\mu_2(x)$ specified, we define component unreliability as

$$P_D(X \in \mathcal{S}_2; \mu_2(x)) \propto \int_x \mu_2(x) \pi_D(x \in \mathcal{S}_2) dP_D(x). \quad (4)$$

Components in Vague Binary States (cont.)

- We could define component unreliability as

$P_D(X \in \mathcal{S}_1^C; \mu_1(x))$, where \mathcal{S}_1^C has membership function, $1 - \mu_1(x)$. In this case, the component unreliability is

$$P_D(X \in \mathcal{S}_1^C; \mu_1(x)) \propto \int_x (1 - \mu_1(x))[1 - \pi_D(x \in \mathcal{S}_1)]dP_D(x). \quad (5)$$

- When a component's state is vague and binary, its unreliability is not the complement of its reliability!

Vague Binary Systems

- Claim: the structure functions of binary state coherent systems with precise classification are membership functions of precise sets
- Let $X_i =$ state of component i , taking values in $\mathcal{S} = \{x; 0 \leq x \leq 1\}$
- $\mathcal{S}_1^{[i]} \subset \mathcal{S}$, where $\mathcal{S}_1^{[i]} = \{x_i; x_i \text{ is a "desirable" state}\}$ with associated membership function, $\mu_1^{[i]}(x)$, $i = 1, \dots, n$
- Consider $\mathcal{S}_1^{[i]} \subset \mathcal{S}$ precise $\forall i$. e.g., $\forall i, \exists x_i \ni$
$$\mu_1^{[i]}(x) = \begin{cases} 1 & x \geq x_i^* \\ 0 & x < x_i^* \end{cases}$$

Vague Binary Systems

- Let $\mathbf{X} = (X_1, \dots, X_n)$ and suppose that the n components are connected in series. Thus $\phi_S(\mathbf{X}) = 1$ if and only if $x \geq x_i^*$ for all i , $i = 1, \dots, n$, implying

$$\phi_S(\mathbf{X}) = \prod_{i=1}^n \mu_1^{[i]}(X_i) = \min_i [\mu_1^{[i]}(X_i)], \quad (6)$$

- Similarly, if n components were connected in parallel redundancy, then

$$\phi_P(\mathbf{X}) = \prod_{i=1}^n \mu_1^{[i]}(X_i) = \max_i [\mu_1^{[i]}(X_i)] \quad (7)$$

- For a k -out-of- n system, $\phi_K(\mathbf{X}) = \mu_1^{[(n-k+1)]}(\mathbf{X})$ where $\mu_1^{[(n-k+1)]}(\mathbf{X})$ represents the $(n - k + 1)$ st membership function when $\mu_1^{[i]}(X_i)$'s are ordered

Vague Binary Systems

Motivated by the above, we define system structure functions as

$$\begin{aligned}\phi_S(\mathbf{X}) &= \min_i[\mu_1^{[i]}(X_i)] = \mu_1^{[(1)]}(\mathbf{X}), \\ \phi_P(\mathbf{X}) &= \max_i[\mu_1^{[i]}(X_i)] = \mu_1^{[(n)]}(\mathbf{X}), \text{ and} \\ \phi_K(\mathbf{X}) &= \mu_1^{[(n-k+1)]}(\mathbf{X})\end{aligned}$$

Vague Binary Systems

There are two different strategies for defining the reliability of a vague coherent system.

1. Assume that a system is reliable if the necessary components are in a desirable state

- The reliability of a series system is $\prod_{i=1}^n [P_D(X_i \in \mathcal{S}_1^{[i]}; \mu_1^{[i]}(x_i))]$ assuming independent X_i 's, where

$$P_D(X_i \in \mathcal{S}_1^{[i]}; \mu_1^{[i]}(x_i)) \propto \int_{x_i} \mu_1^{[i]}(x_i) \pi_D(x_i \in \mathcal{S}_1^{[i]}) dP_D^{[i]}(x_i), \quad (8)$$

- Assuming independent X_i 's, the reliability of a parallel redundant system is

$$P_D\left(\bigcup_{i=1}^n \{X_i \in \mathcal{S}_1^{[i]}\}; \mu_1^{[i]}(x_i), i = 1, \dots, n\right)$$

- The case of k -out-of- n systems follows along similar lines

Vague Binary Systems

2. $\mathcal{S}_{\phi(\mathbf{X})}^* = \{x; \phi(\mathbf{X}) = x \text{ is a "desirable" state of the system}\} \subset \mathcal{S}$
with associated membership function $\mu_{\phi(\mathbf{X})}^*(x)$.

- The reliability of a series system is

$$P_D(\phi_S(\mathbf{X}) \in \mathcal{S}_{\phi_S(\mathbf{X})}^*; \mu_{\phi_S(\mathbf{X})}^*(x)) \propto \int_x \mu_{\phi_S(\mathbf{X})}^*(x) \pi_D(x \in \mathcal{S}_{\phi_S(\mathbf{X})}^*) dP_D(x),$$

where $dP_D(x)$ is obtained from

$$P_D(\phi_S(\mathbf{X}) \geq x) = \prod_{i=1}^n P_D(X_i \geq \mu_1^{[i]-1}(x)),$$

assuming independent X_i 's; $\mu_1^{[i]-1}(x)$ denotes the inverse of $\mu_1^{[i]}(x)$.

Vague Binary Systems (cont.)

- Similarly for parallel systems,

$$P_D(\phi_P(\mathbf{X}) \in \mathcal{S}_{\phi_P(\mathbf{X})}^*; \mu_{\phi_P(\mathbf{X})}^*(x)) \propto \int_x \mu_{\phi_P(\mathbf{X})}^*(x) \pi_D(x \in \mathcal{S}_{\phi_P(\mathbf{X})}^*) dP_D(x),$$

where $dP_D(x)$ is obtained via $\prod_{i=1}^n P_D(X_i \leq \mu_1^{[i]-1}(x))$ assuming independent X_i 's.

- The case of $(n - k + 1)$ -out-of- n follows by considering the distribution of the k th order membership function, $\mu_1^{[(k)]}(\mathbf{X})$.

Ongoing Research

- Reliability of $(m + 1)$ -level degraded systems
- Other applications
 - Quality of life
 - Disclosure limitation

References

- Bellman, R.E. and Giertz, M. (1973) On the analytic formalism of the theory of fuzzy sets. *Information Sciences*, 5: 149-157.
- Bement, T., Booker, J., Sellers, K. and Singpurwalla, N. Membership Functions and Probability Measures of Fuzzy Sets. Technical report LA-UR-00-3660, Los Alamos National Laboratory, Los Alamos, NM. August, 2000.
- Black, M. (1939) Vagueness: an Exercise in Logical Analysis *Philosophy of Science*. 427-455.

References (cont.)

- Lukasiewicz, J. (1930) Philosophische Bemerkungen zu mehrwertigen Systemen des Aussagenkalküls. *Comptes rendus des séances de la Société des Sciences et des Lettres de Varsovie Cl. III*, 23, 51-77; English tr. Philosophical remarks on many-valued systems of propositional logic [in:] Mc-Call, X. (ed.) *Polish Logic 1920-1939*. Clarendon Press, Oxford, 1967, 40-65.
- Malinowski, G. (1993) *Many-valued logics*. Clarendon Press, Oxford.
- Zadeh, L.A. (1965) “Fuzzy Sets”. *Information and Control*, 8, 338-353.
- Zadeh, L.A. (1968) Fuzzy Sets as a Basis for a Theory of Possibility. *Fuzzy Sets and Systems*, 3-28.