

Issued Under Office of Naval Research
Contract No. Nonr- 1858(16)

RETAIN THIS COPY
FOR YOUR FILES
USE JOB NO. BELOW
<input type="radio"/> TO CHECK YOUR BILLS
<input type="radio"/> TO REORDER
<input type="radio"/> FOR ALL REFERENCES
JOB NO. 4069-62
No. Copies
APR 10 1962
PRINCETON UNIVERSITY DUPLICATING BUREAU

n-PERSON GAMES WITH ONLY
1, n-1, AND n-PERSON PERMISSIBLE COALITIONS

Michael Maschler

Econometric Research Program
Research Memorandum No. 37
19 March 1962

The research described in this paper was supported partially by the Office of Naval Research. Reproduction, translation, publication, use and disposal in whole or in part by or for the United States Government is permitted.

Princeton University
Econometric Research Program
92-A Nassau Street
Princeton, N. J.

Maschler

37

1. Introduction.

The purpose of this paper is to present a complete description of various bargaining sets for n -person games, in which the set of permissible coalitions consists of the 1, $n-1$, and n -person coalitions. These games can be regarded as special $(n-1)$ -quota games (see definitions in Section 3), and it turns out that the forms the bargaining sets take depend on whether or not the $(n-1)$ -quota is non-negative.

Theoretically, the bargaining sets can always be computed as solutions of certain systems of linear inequalities in the payoff space, connected by the words "and" and "or." (See R. J. Aumann and M. Maschler, [1].) In fact, the computation of the bargaining set \mathcal{M} in our case was carried out (in [1]) for $n = 3, 4$. In general, however, the computation is very difficult unless some short-cuts can be found. In this paper we used the method of "deleting" "weak players" from the original game, thus being able to obtain a very simple inductive method for constructing the bargaining sets.

It turns out that the various bargaining sets happen to be the same for the games treated here, except that, if the game has an empty core, outcomes of the form $(x_1, x_2, \dots, x_n; 1, 2 \dots n)$ occur only in some of the bargaining sets.

For the sake of completeness we repeat the necessary definitions. These first appeared in [1].

2. Preliminary notations and definitions.

We are concerned with an n -person cooperative game Γ , given by a set $N = \{1, 2, \dots, n\}$ of n players, a set $\{B\}$ of non-empty subsets of N , called permissible coalitions, and a real valued characteristic function $v(B)$ defined for the elements B of $\{B\}$.

For the sake of simplicity and for normalization purposes, we assume that¹

$$(2.1) \quad i \in \{B\}, \quad v(i) = 0 \quad \text{for each } i, \quad i \in N.$$

$$(2.2) \quad v(B) \geq 0 \quad \text{for each } B, \quad B \in \{B\}.$$

An outcome of a game in which the players are partitioned into disjoint permissible coalitions B_1, B_2, \dots, B_m , where each coalition shares its value among its members, will be described by a "payoff configuration" (p.c.):

$$(2.3) \quad (x; \mathcal{B}) \equiv (x_1, x_2, \dots, x_n; B_1, B_2, \dots, B_m).$$

Here,

$$(2.4) \quad B_j \in \{B\}, \quad B_j \cap B_k = \phi \quad \text{for } j \neq k, \quad \bigcup_{j=1}^m B_j = N, \quad j, k = 1, 2, \dots, m,$$

and x_i is a real number which represents the payoff to player i , $i = 1, 2, \dots, n$, and therefore

$$(2.5) \quad \sum_{i \in B_j} x_i = v(B_j).$$

$\mathcal{B} \equiv B_1, B_2, \dots, B_m$ will be called the coalition structure and $x \equiv (x_1, x_2, \dots, x_n)$ will be referred to as the payoff.

Let K be a non-empty set of players. If the players in K want to get their shares in $(x; \mathcal{B})$, they need only the consent of the players in those coalitions which intersect K . The members of these coalitions will be called the partners of K in $(x; \mathcal{B})$, and the set of the partners will be denoted by $P[K; (x; \mathcal{B})]$. Thus

$$(2.6) \quad P[K; (x; \mathcal{B})] \equiv \{i \mid i \in B_j, B_j \cap K \neq \phi\}.$$

¹In Section 4 we shall encounter a case where the normalization $v(i) = 0, i = 1, 2, \dots, n$, will not occur. Nothing in the definitions in this section should be changed if the game is not normalized.

Note that in this terminology, $K \subset P[K; (x; \mathcal{B})]$. If S is a subset of a coalition B , we shall refer to the players in the complement of S with respect to B as the associates of S in B .

We shall be interested in "stable" payoff configurations, which will later be defined. One of the demands which we shall impose on them is that they should be coalitionally rational, in the sense that no subcoalition of an already existing coalition could alone make more. Accordingly, we make the following

Definition 2.1 A p.c. (2.3) for a game Γ will be called coalitionally rational (c.r.p.c.) if

$$(2.7) \quad \sum_{i \in B} x_i \geq v(B) \quad \text{for all } B, B \subset B_j, j = 1, 2, \dots, m, B \in \{B\}.$$

A c.r.p.c. is certainly individually rational, but x in $(x; \mathcal{B})$ does not have to belong to the core of the game, because (2.7) has to be satisfied only for each permissible $B, B \subset B_j, j = 1, 2, \dots, m$.

Definition 2.2 Let $(x; \mathcal{B})$ be a c.r.p.c. (2.3), for a game Γ , and let K and L be two non-empty disjoint subsets of a coalition B_j which occurs in $(x; \mathcal{B})$:

$$(2.8) \quad K, L \neq \phi, K \cap L = \phi, K, L \subset B_j, 1 \leq j \leq m.$$

An objection of K against L in $(x; \mathcal{B})$ is a c.r.p.c.

$$(2.9) \quad (y; \mathcal{C}) \equiv (y_1, y_2, \dots, y_n; C_1, C_2, \dots, C_p),$$

for which

$$(2.10) \quad P[K; (y; \mathcal{C})] \cap L = \phi,$$

$$(2.11) \quad y_i > x_i \quad \text{for all } i, i \in K,$$

$$(2.12) \quad y_i \geq x_i \quad \text{for all } i, i \in P[K; (y; \mathcal{C})].$$

Thus, in an objection $(y; \mathcal{C})$, the players in K claim that they can get more ((2.11)) without the aid of the players L ((2.10)), and the new situation is reasonable, because $(y; \mathcal{C})$ is a c.r.p.c. in which the partners of K get not less than they got in $(x; \mathcal{B})$ ((2.12)).

Definition 2.3 Let $(x; \mathcal{B})$ be a c.r.p.c. (2.3) in a game Γ , and let $(y; \mathcal{C})$ be an objection of a set K against a set L in $(x; \mathcal{B})$. K, L satisfy (2.8). A counter objection of L against K is a c.r.p.c.

$$(2.13) \quad (z; \mathcal{D}) \equiv (z_1, z_2, \dots, z_n; D_1, D_2, \dots, D_q),$$

for which

$$(2.14) \quad P[L; (z; \mathcal{D})] \not\supset K,$$

$$(2.15) \quad z_i \geq x_i \text{ for all } i, i \in P[L; (z; \mathcal{D})],$$

$$(2.16) \quad z_i \geq y_i \text{ for all } i, i \in P[L; (z; \mathcal{D})] \cap P[K; (y; \mathcal{C})].$$

In their counter objection $(z; \mathcal{D})$, the players in L claim that they can protect their share ((2.15)) by giving their partners at least what they had before ((2.15)), and if they need the consent of some of the partners of K who were included in the objection, they can offer them not less than what they were offered in the objection ((2.16)). The members of L are allowed to use the tactics of "divide and rule," by taking some members of K as partners, but they may not take all the members of K as partners ((2.14)).

Definition 2.4 A c.r.p.c. $(x; \mathcal{B})$ in a game Γ is called stable, if for each objection of a set K against a set L in $(x; \mathcal{B})$ there is a counter objection of L against K .

The set \mathcal{M} of all the stable p.c.'s in a game Γ will be called the bargaining set of Γ . This bargaining set was defined and some of its

properties were explored in R. J. Aumann - M. Maschler [1].

3. Effective coalitions.

Definition 3.1 Let B^* be a permissible coalition in a game Γ . B^* will be called an effective coalition if there exists a payoff $\{x_i\}$, $i \in B^*$, such that

$$(3.1) \quad \sum_{i \in B^*} x_i = v(B^*),$$

$$(3.2) \quad \sum_{i \in B} x_i \geq v(B) \text{ for all } B, B \subset B^*, B \in \{B\}.$$

Discussion. A coalition is effective if and only if its value can be shared among its members in such a way that no permissible subcoalition is able alone to make more. Such a share will be called an effective share. It follows from Definition 2.1 that only effective coalitions appear in a c.r.p.c., and the payoff of a c.r.p.c. induces an effective share in each one of its coalitions. Therefore, a non-effective coalition can never enter a stable p.c., nor can it be used for objections or counter objections. Accordingly, the bargaining set of a game Γ will not change if one declares the non-effective coalitions of Γ as not permissible.

Theorem 3.1 Let B^* be a permissible k -person coalition, $k \geq 2$, in a game Γ . Suppose also that any permissible subcoalition of B^* which contains at most $k - 2$ players has a zero value. Let $B^{(v)}$ denote the coalition $B^* - \{v\}$, $\{v\} \in B^*$, and let $v^{(v)} = v(B^{(v)})$ if $B^{(v)}$ is a permissible coalition, and $v^{(v)} = 0$ if $B^{(v)}$ is not permissible.¹ A necessary and sufficient condition for the coalition B^* to be effective is:

$$(3.3) \quad v(B^*) \geq v^{(v)} \text{ for all } v, v \in B^*,$$

$$(3.4) \quad (k-1) v(B^*) \geq \sum_{i \in B^*} v^{(i)}.$$

¹The bargaining set is not changed essentially if we replace the non-permissible coalitions in a game by permissible coalitions having a zero value.

Proof: If B^* is effective, let $\{x_i\}$, $i \in B^*$, be an effective share of its value among its members. Certainly $x_i \geq 0$ for all i , $i \in B^*$. Applying (3.2) to the coalitions $B^{(v)}$, $v \in B^*$, we obtain, in view of (3.1), the condition (3.3). Summing up the same inequalities (3.2) and dividing both sides by $k-1$, we obtain (3.4). The conditions (3.3) and (3.4) are therefore necessary.

If (3.3) and (3.4) hold, it is possible to choose numbers $w^{(v)}$, $v \in B^*$, which satisfy

$$(3.5) \quad (k-1) v(B^*) = \sum_{i \in B^*} w^{(i)},$$

$$(3.6) \quad v(B^*) \geq w^{(v)} \quad \text{for all } v, v \in B^*,$$

$$(3.7) \quad w^{(v)} \geq v^{(v)} \quad \text{for all } v, v \in B^*.$$

Indeed, we increase continuously the $v^{(v)}$'s one by one so as not to exceed $v(B^*)$, until the sum of the new $v^{(v)}$'s becomes equal to $(k-1) v(B^*)$. We then denote these new $v^{(v)}$'s by $w^{(v)}$'s. We shall certainly reach the required sums because of (3.4) and because $k v(B^*) \geq (k-1) v(B^*)$. We now choose the payoffs $\{x_v\}$ to be:

$$(3.8) \quad x_v = v(B^*) - w^{(v)}, \quad v \in B^*.$$

By (3.6), $x_v \geq 0$ for all v , $v \in B^*$. Also, by (3.8), (3.5) and (3.6), we get for each v , $v \in B^*$,

$$(3.9) \quad \sum_{i \in B^{(v)}} x_i = (k-1) v(B^*) - \sum_{i \in B^{(v)}} w^{(i)} = w^{(v)} \geq v^{(v)}.$$

In addition, by (3.8) and (3.5),

$$(3.10) \quad \sum_{i \in B^*} x_i = k v(B^*) - \sum_{i \in B^*} w^{(i)} = v(B^*).$$

This shows that $\{x_v\}$, $v \in B^*$, is an effective share of $v(B^*)$ among the members of B^* , hence B^* is effective. The conditions (3.3) and (3.4) are therefore sufficient.

Corollary 3.1 It is easy to check that if

$$(3.11) \quad (k-1) v^{(v)} \leq \sum_{i \in B^*} v^{(i)} \quad \text{for all } v, v \in B^*,$$

then condition (3.4) implies condition (3.3). If, however, condition (3.11) is not satisfied, then condition (3.3) implies condition (3.4).

Corollary 3.2 We shall call the conditions (3.11) generalized triangle inequalities, because of the following interesting properties:

First, they reduce to the triangle inequalities if $k = 3$.

Secondly, if these conditions are satisfied, and B^{**} is any subset of B^* , containing r members, then the numbers $v^{(i)}$, $i \in B^{**}$, also satisfy the generalized triangle inequalities. (In particular, any three distinct numbers among the $v^{(v)}$'s satisfy the triangle inequalities.)

Proof: By finite induction, it is sufficient to prove that any $k-1$ distinct numbers among the $v^{(v)}$'s satisfy the generalized triangle inequalities. Let B^{**} be a subset of B^* containing $k-1$ elements, and let v_0 belong to $B^* - B^{**}$; i.e., $B^{**} = B \setminus \{v_0\}$. We may write the inequality (3.11), for $v = v_0$, in the form

$$(3.12) \quad (k-2) v^{(v_0)} \leq \sum_{i \in B} v^{(i)}.$$

Multiplying the remaining inequalities of (3.11) by $(k-2)$ and substituting (3.12) for $(k-2) v^{(v_0)}$, we obtain

$$(3.13) \quad (k-1)(k-2) v^{(v)} \leq (k-2) \sum_{i \in B} v^{(i)} + \sum_{i \in B} v^{(i)}$$

for all $v, v \neq v_0, v \in B^*$. Therefore,

$$(3.14) \quad (k-2) v^{(v)} \leq \sum_{i \in B^{**}} v^{(i)} \quad \text{for all } v, v \in B^{**}.$$

Corollary 3.3 If the generalized triangle inequalities (3.11) do not all hold, and if

$$(3.15) \quad v^{(v_0)} = \text{Max}_{v \in B^*} v^{(v)},$$

then

$$(3.16) \quad (k-1) v^{(v_0)} > \sum_{i \in B^*} v^{(i)}.$$

Proof: If

$$(3.17) \quad (k-1) v^{(v_1)} > \sum_{i \in B^*} v^{(i)}$$

and

$$(k-1) v^{(v_2)} \leq \sum_{i \in B^*} v^{(i)},$$

then certainly

$$v^{(v_1)} > v^{(v_2)}.$$

4. n-Person, (n-1)-quota games.

Following L. S. Shapley [4], G. K. Kalisch [2], we introduce the following

Definition 4.1 An n-person game Γ , $n \geq 3$, is an m-quota game, if $2 \leq m < n$, if all the m-person coalitions are permissible and if there exist real numbers $\omega_1, \omega_2, \dots, \omega_n$ such that^{1,2}

$$(4.1) \quad v(B) = \sum_{i \in B} \omega_i \quad \text{for all } B, |B| = m.$$

The vector $(\omega_1, \omega_2, \dots, \omega_n)$ will be called the m-quota (or simply the quota) of the game Γ , and the number ω_i will be referred to as the quota of player i , $i = 1, 2, \dots, n$. A player whose quota is negative (where the game is normalized by (2.1)) will be called a weak player. If the order of the quota is unclear from the contents, we shall say that

¹ $|B|$ denotes the number of players in B .

² Our definition is somewhat more general than the definitions given in [2], [4].

the m-quota "contains" a weak player, or that a certain player is weak in the game with respect to the m-quota.

Lemma 4.1 There are at most m-1 weak players in an m-quota game.

Proof: This follows immediately from (2.2) and (4.1).

Lemma 4.2 The m-quota of a game, if such exists, is unique.

Proof: Choose an arbitrary (m+1)-person subset E of N . (E exists because $m < n$). Consider the $m + 1$ m-person subsets of E and apply to them the equations (4.1). The resulting linear equations in the ω_i 's , $i \in E$, will have a unique solution. Thus, any $m + 1$ ω_i 's are uniquely determined, hence the quota is unique.

Lemma 4.3 An n-person game, $n \geq 3$, in which the (n-1)-person coalitions are permissible, is an (n-1)-quota game. The quota is

$$(4.2) \quad \omega_v = \frac{v^{(1)} + v^{(2)} + \dots + v^{(n)} - (n-1)v^{(v)}}{n - 1} , \quad v = 1, 2, \dots, n ,$$

where

$$(4.3) \quad v^{(i)} \equiv v(N^{(i)}) , \quad N^{(i)} \equiv N - \{i\} , \quad i = 1, 2, \dots, n .$$

Proof: Equations (4.1) are satisfied.

Lemma 4.4 An n-person game, regarded as an (n-1)-quota game, has no weak player, if and only if the values $v^{(i)}$, $i = 1, 2, \dots, n$, defined by (4.3), satisfy the generalized triangle inequalities:

$$(4.4) \quad (n-1) v^{(v)} \leq v^{(1)} + v^{(2)} + \dots + v^{(n)} , \quad v = 1, 2, \dots, n .$$

(See Corollary 3.2.)

The proof follows immediately from (4.2).

In this section we shall analyze n-person games in which the sets of the permissible coalitions consist of the 1, n-1, and n-person coalitions. We shall characterize completely the bargaining sets for these games.

Theorem 4.1 Let Γ be an n-person game in which the set of the permissible coalitions consists of the 1, n-1, and n-person coalitions. A p.c. of the form¹

$$(4.5) \quad (\mathbf{x}; N) \equiv (x_1, x_2, \dots, x_n; N)$$

is stable if and only if N is an effective coalition and \mathbf{x} is an effective share of $v(N)$ among the players.

Proof: A necessary and sufficient condition that $(\mathbf{x}; N)$ is a c.r.p.c. is that N is effective and \mathbf{x} is an effective share of $v(N)$ among the players. A necessary condition that $(\mathbf{x}; N)$ is stable is that it is a c.r.p.c. This condition is also sufficient, because if $(\mathbf{x}; N)$ is a c.r.p.c., no objections are possible.

Corollary 4.1 If $n \geq 2$, then Theorem 3.1 gives a necessary and sufficient condition for N to be effective, namely:

$$(4.6) \quad (n-1) v(N) \geq \sum_{i=1}^n v^{(i)}, \quad v(N) \geq v^{(i)}, \quad i = 1, 2, \dots, n,$$

where $v^{(i)}$ is defined by (4.3).

The value of $v(N)$ and the fact that it is permissible or not have no effect on the stability or instability of p.c.'s of the form $(\mathbf{x}; \mathcal{B})$, where $\mathcal{B} \neq N$, because by (2.10) and (2.14), the coalition N cannot be formed in any objection or any counter objection.

Obviously, the p.c. $(0, 0, \dots, 0; 1, 2, \dots, n)$ is always stable. We have yet to find the stable p.c.'s of the form $(\mathbf{x}; N - \{j\}, j)$.

¹"of the form" will mean in this paper "having the chosen coalition structure."

It will turn out that the occurrence or non-occurrence of a weak player in the $(n-1)$ -quota plays an important role in the result.

Theorem 4.2 Let Γ be an n -person game, $n \geq 3$, in which the set of the permissible coalitions consists of the 1 , $n-1$, and n -person coalitions.

Let $\omega_1, \omega_2, \dots, \omega_n$ be the $(n-1)$ -quota of Γ . If no player is weak, then a p.c. of the form $(x; j, N^{(j)})$, where $N^{(j)} \equiv N - \{j\}$, is stable if and only if

$$(4.7) \quad (x; j, N^{(j)}) = (\omega_1, \omega_2, \dots, \omega_{j-1}, 0, \omega_{j+1}, \dots, \omega_n; j, N^{(j)}) .$$

Remark. The quota is given by (4.2). Necessary and sufficient conditions that no player is weak are given in Lemma 4.4. This theorem asserts that if the numbers $v(N^{(j)})$, $j = 1, 2, \dots, n$, satisfy the generalized triangle inequalities, then the stable p.c.'s are such that each player gets his quota if he is a member of an $(n-1)$ -person coalition.

Proof: Being an $(n-1)$ -quota, the ω_i 's satisfy

$$(4.8) \quad \omega_1 + \omega_2 + \dots + \omega_{j-1} + \omega_{j+1} + \dots + \omega_n = v^{(j)}, \quad j = 1, 2, \dots, n ,$$

where $v^{(j)} \equiv v(N^{(j)})$. The quota for each player is non-negative, hence

(4.7) is a c.r.p.c. To show that it is stable, we observe that in any possible objection in (4.7), if any¹, a coalition $N^{(\ell)}$ is formed, $\ell \neq j$.

Such an objection is an objection of a set K against player ℓ , where K is a non-empty subset of some players, each different from player j , who receive in the objection more than their quotas. The objection must therefore be of the form:

$$(4.9) \quad (y; \ell, N^{(\ell)}) = (\omega_1 + \alpha_1, \omega_2 + \alpha_2, \dots, \omega_{\ell-1} + \alpha_{\ell-1}, 0, \omega_{\ell+1}, \dots, \omega_n + \alpha_n; \ell, N^{(\ell)}) ,$$

¹No objection is possible if $\omega_j = 0$.

where $\alpha_i \geq 0$ for $i \neq \ell, j$, $\alpha_i > 0$ for $i \in K$, $\ell, j \notin K$, $\omega_j + \alpha_j \geq 0$ and $\alpha_1 + \alpha_2 + \dots + \alpha_{\ell-1} + \alpha_{\ell+1} + \dots + \alpha_n = 0$. (The last inequality follows from (4.8), where j is replaced by ℓ .) Thus $\alpha_j < 0$.

Let μ be a player in the objecting K ; then $\alpha_\mu > 0$. A counter objection can now be formed as follows¹:

$$(4.10) \quad (z; \mu, N^{(\mu)}) = (\omega_1 + \alpha_1, \omega_2 + \alpha_2, \dots, \omega_{\ell-1} + \alpha_{\ell-1}, \omega_\ell + \alpha_\mu, \\ \omega_{\ell+1} + \alpha_{\ell+1}, \dots, \omega_{\mu-1} + \alpha_{\mu-1}, 0, \omega_{\mu+1} + \alpha_{\mu+1}, \dots, \omega_n + \alpha_n; \mu, N^{(\mu)}).$$

Evidently, $z \geq 0$ coordinatewise. Also, the sum of its coordinates is $v^{(\mu)}$ (see (4.8)). Hence (4.10) is a c.r.p.c. It is a counter objection because $\alpha_\mu > 0$. We have proved that (4.7) is stable.

Suppose now that

$$(4.11) \quad (x; \mathcal{B}) \equiv (x_1, x_2, \dots, x_{j-1}, 0, x_{j+1}, \dots, x_n; j, N^{(j)}) \in \mathcal{M},$$

and suppose that $x_\mu < \omega_\mu$, $\mu \neq j$. Then, there exists at least one player v , such that $x_v > \omega_v$. Let ϵ be a positive number, smaller than $x_v - \omega_v$, then player μ can object against player v by²

$$(4.12) \quad (x_1, x_2, \dots, x_{\mu-1}, x_\mu + \epsilon, x_{\mu+1}, \dots, x_{j-1}, \bar{y}_j, x_{j+1}, \dots, \\ x_{v-1}, 0, x_{v+1}, \dots, x_n; v, N^{(v)}).$$

Here, by (4.8),

$$(4.13) \quad \bar{y}_j = v^{(v)} - \sum_{i \neq j, v} x_i - \epsilon = (\omega_1 + \omega_2 + \dots + \omega_n - \omega_v) - \\ - (\omega_1 + \omega_2 + \dots + \omega_n - \omega_j - x_v) - \epsilon = \omega_j + x_v - \omega_v - \epsilon > \omega_j \geq 0.$$

Therefore, (4.13) is an objection.

Player v has no counter objection. Indeed, he cannot counter object by

¹Despite the notation in (4.10), we do not assume $\mu > \ell$. However, $\mu \neq \ell$.

²Despite the notation in (4.12), we assume no order relation among the numbers μ, v, j , except that they are distinct.

playing as a 1-person coalition, because $x_v > \omega_v \geq 0$. He cannot appeal to the coalition N because of the restriction (2.14). By (2.14) he can only try to object by participating in the coalition $N^{(\mu)}$; but, summing up the minimum amounts that the members of this coalition must be paid in a counter objection, we obtain by (4.8), (4.11), and (4.13):

$$\begin{aligned} & x_1 + x_2 + \dots + x_{\mu-1} + x_{\mu+1} + \dots + x_{j-1} + y_j + x_{j+1} + \dots + x_{v-1} + x_v + \\ & + x_{v+1} + \dots + x_n = (\omega_1 + \omega_2 + \dots + \omega_n - \omega_j - x_\mu) + (\omega_j + x_v - \omega_v - \epsilon) > \\ & > (\omega_1 + \omega_2 + \dots + \omega_n - \omega_\mu) + (x_v - \omega_v - \epsilon) = v^{(\mu)} + (x_v - \omega_v - \epsilon) > v^{(\mu)}. \end{aligned}$$

This sum is greater than the amount the coalition $N^{(\mu)}$ can make, hence a counter objection is not possible.

We have proved that $(x; \mathcal{B})$ is not stable, contrary to the assumption (4.11), hence each player in $(x; \mathcal{B})$ must receive at least, and therefore exactly, his quota. This completes the proof of the theorem.

If a weak player is present in the $(n-1)$ -quota, we shall see that the game is essentially reduced to a similar $(n-1)$ -person game. To this effect, we shall now explain how to "delete" players from our games.

Definition 4.2. Let Γ be an n -person game, $n \geq 3$, in which the set of the permissible coalitions consists of the 1, $n-1$, and n -person coalitions. A game Γ_v is said to result from game Γ by deleting player v , if:

- (i) The players in Γ_v form the set $N - \{v\}$.
- (ii) The set of permissible coalitions in Γ_v consists of the 1, $n-2$, and $(n-1)$ -person coalitions.
- (iii) The characteristic function $v_v(B)$ of Γ_v satisfies

$$(4.14) \quad v_v(B^{(v)}) = v_v(B),$$

whenever B is an $(n-1)$ -person coalition in Γ and $B^{(v)} \equiv B - \{v\}$, and, if $n > 3$,

$$(4.15) \quad v_v(i) = 0 \text{ for all } i, i \in N - \{v\} .$$

Discussion: Note that $B^{(v)} = B$ if $v \notin B$. In deleting player v from the game Γ , we ignored the n -person coalition, and from each $(n-1)$ -person coalition which contained player v we made an $(n-2)$ -person coalition by removing this player from the coalition without altering the value of the coalition. The $(n-1)$ -person coalition which did not include player v remained unchanged. If $n = 3$, it may well happen that the characteristic function does not satisfy the normalization (2.1).

Lemma 4.5 If $\{\omega_i\}$, $i = 1, 2, \dots, n$ is the $(n-1)$ -quota of Γ , then $\{\omega_i + \omega_v / (n-2)\}$, $i = 1, 2, \dots, v-1, v+1, \dots, n$ is the $(n-2)$ -quota of Γ_v .

The proof is immediate.

Corollary 4.2 If we order the players by their quotas, then deleting a player from the game will not change the order relation for the rest of the players. In particular, if we delete a weak player, the remaining weak players (if such exist) will remain weak, some previously non-weak players may now become weak, but their new quotas will be greater than the new quotas of the previously weak players.

Lemma 4.6 Let Γ be an n -person game, in which the set of the permissible coalitions consists of the $1, n-1,$ and n -person coalitions. Let

$$(4.16) \quad (\mathbf{x}; j, N^{(j)}) \equiv (x_1, x_2, \dots, x_{j-1}, 0, x_{j+1}, \dots, x_n; j, N - \{j\})$$

be a c.r.p.c. in Γ . Then:

(i) An objection of the form

$$(4.17) \quad (\mathbf{y}; \ell, N^{(\ell)}) \equiv (y_1, y_2, \dots, y_{\ell-1}, 0, y_{\ell+1}, \dots, y_n; \ell, N - \{\ell\})$$

exists¹ if and only if $\ell \neq j$ and

¹By (2.10) and (2.11), any possible objection must be of this form.

$$(4.18) \quad \sum_{i \neq j, \ell} x_i < v^{(\ell)} .$$

(ii) If an objection of the form (4.17) exists, for a fixed ℓ , $\ell \neq j$, then each objection of this form can be countered, if and only if

$$(4.19) \quad x_\ell = 0 \text{ or } v^{(k)} + x_k \geq v^{(\ell)} + x_\ell \text{ for all } k, k \neq j .$$

Here $v^{(v)} \equiv v(N^{(v)}) \equiv v(N - \{v\})$, $v = 1, 2, \dots, n$.

Proof: The necessity and sufficiency of condition (4.18) follows immediately from Definition 2.2. Suppose that (4.18) is satisfied, and that any objection of the form (4.17) can be countered, for a fixed ℓ . Each objection of this form must be regarded as an objection of a set K against player ℓ , where K is a non-empty subset of the set consisting of those players, different from player j , for which $y_i > x_i$. In particular, for each fixed player k , $k \in N - \{\ell\} - \{j\}$, we can distribute the amount $v^{(\ell)}$ in such a way that $y_k = x_k + \epsilon$, where ϵ is an arbitrarily chosen small enough positive number, and make this distribution an objection of player k against player ℓ . Player ℓ can counter object in at most two ways: either by playing as a 1-person coalition, which is possible if and only if $x_\ell = 0$; or by joining the coalition $N^{(k)}$, which is possible if and only if

$$(4.20) \quad v^{(k)} - x_\ell \geq v^{(\ell)} - x_k - \epsilon .$$

Thus, either $x_\ell = 0$ must hold or (4.20), for each small enough positive ϵ . We have therefore proved that (4.19) is a necessary condition.

Conversely, if $x_\ell = 0$, player ℓ can always counter object by playing as a 1-person coalition. If (4.19) is satisfied, but $x_\ell > 0$, let $(y; \ell, N^{(\ell)})$ be any objection of K against ℓ . Certainly, K is a non-empty subset of $N - \{\ell\} - \{j\}$. Let k be a player in K , then $y_K > x_k$. We claim that player ℓ has a counter objection against K , of the form¹

¹Despite the notation in (4.21), ℓ is not necessarily greater than k .

$$(4.21) \quad (z; k, N^{(k)}) \equiv (y_1, y_2, \dots, y_{k-1}, 0, y_{k+1}, \dots, y_{\ell-1}, z_\ell, y_{\ell+1}, \dots, y_n; k, N^{(k)}),$$

where

$$(4.22) \quad z_\ell = v^{(k)} - \sum_{i \neq k, \ell} y_i.$$

Indeed, because of (4.19),

$$(4.23) \quad z_\ell = v^{(k)} - v^{(\ell)} + y_k > v^{(k)} - v^{(\ell)} + x_k \geq x_\ell.$$

This completes the proof.

Theorem 4.3 Let Γ be an n-person game, $n \geq 3$, in which the set of the permissible coalitions consists of the 1, n-1, and n-person coalitions.

Suppose that the (n-1)-quota of the game contains a weak player, and let v be the player having the smallest quota. Let Γ_v be the (n-1)-person game which results by deleting player v from Γ . Under these conditions:

(i) A p.c. of the form

$$(4.24) \quad (x; v, N^{(v)}) \equiv (x_1, x_2, \dots, x_{v-1}, 0, x_{v+1}, \dots, x_n; v, N - \{v\})$$

is stable in Γ , if and only if

$$(4.25) \quad (x^*; N^{(v)}) \equiv (x_1, x_2, \dots, x_{v-1}, x_{v+1}, \dots, x_n; N - \{v\})$$

is stable in Γ_v . Such stable p.c.'s always exist.

(ii) A p.c. of the form

$$(4.26) \quad (x; j, N^{(j)}) \equiv (x_1, x_2, \dots, x_{j-1}, 0, x_{j+1}, \dots, x_n; j, N - \{j\}),$$

$j \neq v,$

is stable in Γ if and only if¹ $x_v = 0$ and

$$(4.27) \quad (x^*; j, N^{(v,j)}) \equiv (x_1, x_2, \dots, x_{v-1}, x_{v+1}, \dots, x_{j-1}, 0, x_{j+1}, \dots, x_n; j, N - \{v\} - \{j\})$$

¹Despite the notation in (4.27), v is not necessarily smaller than j .

is stable in Γ_v .

Proof: (i) Player v has the smallest quota; hence, by (4.1) and (4.14),

$$(4.28) \quad v_v \equiv v^{(v)} = \text{Max}_{i \in N} v^{(i)} \geq \text{Max}_{i \in N - \{v\}} v^{(i)} = \text{Max}_{i \in N - \{v\}} v_v^{(i)},$$

where the following notation is used for the characteristic function $v_v(B)$ of Γ_v :

$$(4.29) \quad v_v \equiv v_v(N - \{v\}), \quad v_v^{(i)} \equiv v_v(N - \{v\} - \{i\}) \quad \text{for } i \neq v.$$

By corollary 3.3,

$$(4.30) \quad (n-1)v_v = (n-1)v^{(v)} > \sum_{i \in N} v^{(i)} = v_v + \sum_{i \in N - \{v\}} v_v^{(i)}.$$

Therefore,

$$(4.31) \quad (n-2)v_v > \sum_{i \in N - \{v\}} v_v^{(i)}.$$

Corollary 4.1, applied to the game Γ_v , (4.28) and (4.31) insure the effectiveness¹ of the coalition $N - \{v\}$ in Γ_v . Thus, by Theorem 4.1, stable p.c.'s for the game Γ_v , of the form (4.25) always exist.

If (4.25) is stable in the game Γ_v , then certainly (4.24) is a c.r.p.c., because $v_v = v^{(v)}$, and because $x^* \geq 0$ coordinatewise. We know already that 1 and n -person coalitions cannot be used in any objection to (4.24). The p.c. (4.25), being stable in Γ_v , induces by Theorem 4.1 an effective share of v_v among the players of $N - \{v\}$. This means that

$$(4.32) \quad \sum_{i \neq v, \ell} x_i \geq v_v^{(\ell)} \equiv v^{(\ell)} \quad \text{for each } \ell, \ell \neq v.$$

Lemma 4.6, part (i), now implies that no objections are possible in (4.24), with respect to the game Γ . Therefore (4.24) is stable in Γ .

Conversely, if (4.25) is not stable in Γ_v , but (4.24) is stable in Γ , then, by Theorem 4.1, x^* is not an effective share² of v_v among

¹Even if $n = 3$, where (2.1) does not necessarily hold for Γ_v .

²Even in the case $n = 3$, where the assumption (2.1) does not necessarily hold for Γ_v .

the players in Γ_v . Therefore, there exists a coalition $N^{(v,\ell)}$, $v \neq \ell$, for which

$$(4.33) \quad \sum_{i \neq v, \ell} x_i < v_v^{(\ell)} = \bar{v}^{(\ell)}.$$

Lemma 4.6, part (i), now asserts that there exists an objection of the form (4.17) in (4.24), with respect to the game Γ . If we show that (4.19) is not satisfied, then, by Lemma 4.6, part (ii), (4.25) would not be stable in Γ , whence a contradiction. This is indeed the case, because if $x_\ell = 0$, then (4.33) would imply $v_v < v^{(\ell)}$, contradicting (4.28). If for each k , $k \neq v$, $v^{(k)} + x_k \geq v^{(\ell)} + x_\ell$, then (4.33) would imply

$$(4.34) \quad v^{(k)} + x_k > \sum_{i \neq v, \ell} x_i + x_\ell = \sum_{i \neq v} x_i = v^{(v)}.$$

Summing up these inequalities for $k = 1, 2, \dots, v-1, v+1, \dots, n$, we obtain

$$(4.35) \quad \sum_{i=1}^n v^{(i)} > (n-1)v^{(v)}.$$

This contradicts (4.30).

(ii) Consider the expressions (4.26), with $x_v = 0$, and (4.27), with respect to the games Γ and Γ_v , respectively. Certainly, if one of them is a p.c. in its game, so is the other. Also, they are simultaneously coalitionally or non-coalitionally rational in their respective games.¹ We can therefore assume that (4.26) and (4.27) are both c.r.p.c. in their respective games. Applying Lemma 4.6 to the game Γ , we find that the p.c. (4.26) with $x_v = 0$ is stable in Γ if and only if for each ℓ , $\ell \neq j$,

$$(4.36) \quad \sum_{i \neq j, \ell} x_i \geq v^{(\ell)} \quad \text{or} \quad x_\ell = 0 \quad \text{or} \quad v^{(k)} + x_k \geq v^{(\ell)} + x_\ell \quad \text{for all } k, k \neq j.$$

Applying the same Lemma to game Γ_v , and taking (4.14) into account, we find that the p.c. (4.27) is stable in Γ_v if and only if for each ℓ , $\ell \neq j, v$,

¹One has to check especially the case $n = 3$, when the assumption (2.1) does not necessarily hold for Γ_v .

$$(4.37) \quad \sum_{i \neq j, \ell, v} x_i \geq v^{(\ell)} \quad \text{or} \quad x_\ell = 0 \quad \text{or} \quad v^{(k)} + x_k \geq v^{(\ell)} + x_\ell \quad \text{for all } k, \\ k \neq j, v.$$

Certainly, (4.36) implies (4.37), because $x_v = 0$. We shall show that the converse is also true! Indeed, (4.36) is always satisfied for $\ell = v$. Thus, if (4.37) is satisfied but (4.36) does not hold, then a player ℓ , $\ell \neq j, v$ exists, for which

$$(4.38) \quad \sum_{i \neq j, \ell} x_i < v^{(\ell)}, \quad x_\ell > 0, \quad \text{and} \quad v^{(v)} + x_v < v^{(\ell)} + x_\ell.$$

Using again the fact that $x_v = 0$, we find that

$$(4.39) \quad v^{(v)} < v^{(\ell)} + x_\ell \leq v^{(k)} + x_k \quad \text{for all } k, k \neq j, v.$$

Summing up these inequalities for $k \neq j, v$, we obtain

$$(4.40) \quad (n-2)v^{(v)} < \sum_{i \neq j, v} v^{(i)} + \sum_{i \neq j} x_i = \sum_{i \neq v} v^{(i)},$$

or

$$(4.41) \quad (n-1)v^{(v)} < \sum_{i=1}^n v^{(i)}.$$

This contradicts (4.30).

We have proved that the p.c.'s (4.26) with $x_v = 0$, and (4.27) are simultaneously stable or unstable in their respective games. It remains to be shown that the p.c. (4.26) is never stable if¹ $x_v > 0$. We shall show that then (4.36) is not satisfied for $\ell = v$.

It follows from (4.28) that

$$(4.42) \quad \sum_{i \neq j, v} x_i = v^{(j)} - x_v < v^{(j)} \leq v^{(v)}.$$

If

$$(4.43) \quad v^{(k)} + x_k \geq v^{(v)} + x_v \quad \text{for all } k, k \neq j,$$

¹It is certainly unstable if $x_v < 0$, because then (4.26) is not c.r.

then, replacing x_v in (4.43) by 0 and summing these inequalities, we obtain

$$(4.44) \quad \sum_{i \neq j} v^{(i)} + v^{(j)} = \sum_{i=1}^n v^{(i)} > (n-1)v^{(v)} .$$

This contradicts (4.30). Thus, (4.26) cannot be stable if $x_v > 0$, which brings the proof of the theorem to its end.

Conclusion. The procedure for obtaining the bargaining set for the games described in this section is set by Theorems 4.1, 4.2, and 4.3, as follows:

The p.c. $(0, 0, \dots, 0; 1, 2, \dots, n)$ is always stable.

Theorem 4.1 and Corollary 4.1 provide the stable p.c.'s of the form $(x; N)$. These exist if and only if the game has a non-empty core, and the payoffs in the stable p.c.'s form exactly the core of the game.

If the generalized triangle inequalities (4.4) hold for the values of the $(n-1)$ -person coalition, i.e., if no weak player exists in the $(n-1)$ -quota, then Theorem 4.2 provides us with all the stable p.c.'s of the form $(x; j, N-\{j\})$.

If there is a weak player in the game, we delete the weakest player v (i.e., the player with the smallest quota) and reduce the game to an $(n-1)$ -person game with the 1, $n-2$, and $(n-1)$ -person coalitions as permissible. The new game has a non-empty core. The stable p.c. of the form $(x^*; j, N-\{j\}-\{v\})$ in the new game are in 1-1 correspondence to those stable in the original game, of the form $(x; j, N)$. To obtain a stable p.c. in the original game, we pick a stable p.c. in the reduced game, we add the weak player with a 0 payoff, and change the coalition structure in an obvious way.

If, by deleting the weakest players, one at a time, we finally arrive at an m -person game, $m \geq 3$, with no weak player in the $(m-1)$ -quota, we calculate by Theorems 4.1 and 4.2 its bargaining set, from which we can

successively return to the original game, each time by using Theorem 4.1.

If we arrive at a 3-person game, which still contains a weak player, then reducing once more we arrive at a 2-person game of the players, say, k and l , with the characteristic function

$$(4.45) \quad v(k) = a, v(l) = b, v(kl) = c, \quad c > a + b.$$

Obviously, the bargaining set for this game consists of the p.c.'s:

$$(4.46) \quad (a, b; k, l), (a + \alpha, b + \beta; kl), \quad \alpha, \beta \geq 0, \alpha + \beta = c - a - b.$$

From this bargaining set we go back, as before, to the stable p.c.'s of the form $(x; j, N - \{j\})$ in the original game.

Two features should be stressed:

(1) For any coalition-structure of the form $j, N - \{j\}$, there always corresponds a payoff x such that $(x; j, N - \{j\})$ is stable. The game holds true for the coalition-structure N , if and only if N is an effective coalition.

(2) The game has a discrete bargaining set if and only if its $(n-1)$ -quota contains no weak player and the n -person coalition is either ineffective or its value is exactly equal to the sum of the quotas of the players.

5. Other kinds of bargaining sets.

The bargaining set \mathcal{M} represents possible outcomes of a game, assuming that the players wish to end up with that kind of stability which is implied by the definition of \mathcal{M} . Such a wish may sometimes cause inconvenience to the players: It may happen, for example, that the n -person coalition is not effective, yet its value is greater than the value of any other coalition.¹ One may argue that the players might be willing to relax

¹This happens, e.g., in essential constant-sum games (with a superadditive characteristic function).

their stability requirements somewhat, in order to take advantage of forming the n-person coalition.

Taking another point of view, one may argue that an outcome in \mathcal{M} is only protected against threats within an existing coalition, but not against objections raised by members in several existing coalitions against other players in these same coalitions.

In R. J. Aumann and M. Maschler [1], several possible modifications of the theory are suggested for coping with such situations. In this section, we shall examine how some of these modifications affect the bargaining set for the games which were treated in the previous section.

Definition 5.1 A c.r.p.c. $(x; \mathcal{B})$ will be called \mathcal{M}_0 -stable, if it satisfies Definition 2.4, where objections and counter objections are given by Definitions 2.2 and 2.3, except that (2.8) is now replaced by

$$(5.1) \quad K, L \neq \phi, K \cap L = \phi, K \cap B_j \neq \phi \iff L \cap B_j = \phi, \text{ for } B_j \in \mathcal{B}.$$

Verbally, we are now dropping the restriction that K and L belong to one coalition B_j , requiring instead that K and L intersect the same coalitions in \mathcal{B} .

The set of all the \mathcal{M}_0 -stable p.c.'s will be called the bargaining set \mathcal{M}_0 .

Certainly, $\mathcal{M}_0 \subset \mathcal{M}$, and this inclusion may indeed be strict (see [1], [3]).

In this section, in order to be specific, we shall refer to p.c.'s in \mathcal{M} as \mathcal{M} -stable p.c.'s.

Theorem 5.1 Let Γ be an n-person game, in which the set of the permissible coalitions consists of the 1, n-1, and n-person coalitions. The bargaining sets \mathcal{M} and \mathcal{M}_0 for Γ are equal.

Proof: In any possible coalition-structure \mathcal{B} , there occurs at most one coalition which contains more than one player. By (5.1), neither K nor L may contain a 1-person coalition of \mathcal{B} , whence, in our games, condition (5.1) is equivalent to condition (2.8).

Definition 5.2 A c.r.p.c. $(x; \mathcal{B})$ will be called \mathcal{M}_1 -stable, if for each objection of a set K against a set L , there is at least one player in L who can counter-object. (Objections and counter objections are being taken as in¹ Definitions 2.2 and 2.3.)

The set of all the \mathcal{M}_1 -stable p.c.'s will be called the bargaining set \mathcal{M}_1 .

Since an objection of a set K against a set L may be regarded also as an objection of K against any particular member of L , the condition for \mathcal{M}_1 -stability is equivalent to requiring that each player in L is able himself to counter-object (although the players in L may find it impossible to counter-object collectively). Thus, $\mathcal{M} \subset \mathcal{M}_1$, and this inclusion may indeed be strict (see [1]). We further note that any objection of a set of players K against a single player ℓ may be regarded as an objection of a single player k in K , against player ℓ . In view of (2.14), one observes that Definition 5.2 is equivalent to Definition 2.4, when restricting K and L , in the latter case, to containing only one player.

Theorem 5.2 Let Γ be an n-person game, in which the set of the permissible coalitions consists of the 1, n-1, and n-person coalitions. The bargaining sets \mathcal{M} and \mathcal{M}_1 are the same for Γ .

Proof: It is sufficient to prove that $\mathcal{M}_1 \subset \mathcal{M}$ for our games.

¹Although Theorem 5.2 will remain valid if we replace (2.8) by (5.1).

Certainly, there is only one p.c. with the coalition structure $1, 2, \dots, n$, and it is M_1 as well as M -stable.

If $(x_1, x_2, \dots, x_n; N)$ is M_1 -stable, then it is a fortiori a c.r.p.c., and hence, by Theorem 4.1, it is also M -stable.

Suppose that $(x; j, N^{(j)}) \equiv (x_1, x_2, \dots, x_n; j, N - \{j\})$ is M_1 -stable. Let $(y; \mathcal{C})$ be an objection of a set K against a set L , in $(x; j, N^{(j)})$. Clearly, $j \notin K, L$ and therefore \mathcal{C} must be of the form $(\ell, N^{(\ell)})$, $\ell \neq j$. Thus L consists of one player, namely player ℓ , and by M_1 -stability, the objection can indeed be countered.

Definitions 5.3, 5.4, 5.5 The bargaining sets $M^{(i)}$, $M_0^{(i)}$ and $M_1^{(i)}$ are defined exactly as M , M_0 , and M_1 , respectively, except that the coalitional rationality restriction imposed on the p.c.'s is replaced everywhere by the requirement that the p.c.'s are individually rational. I.e., we replace (2.7) by the weaker requirement¹:

$$(5.2) \quad x_i \geq 0 \quad \text{for all } i, i = 1, 2, \dots, n.$$

Clearly, $M_0^{(i)} \subset M^{(i)} \subset M_1^{(i)}$. An example in² [3] shows that the first inclusion may indeed be strict. The following example will show that also the second inclusion may be strict.

Example 5.1³ $n = 4$, $\{B\} = \{i, ij, 123\}$, $v(i) = 0$, $v(ij) = 10$, $v(123) = 18$ for all i, j where $i, j = 1, 2, 3, 4$ and $i \neq j$. Clearly, $(6, 6, 6, 0; 123, 4)$ belongs to $M_1^{(i)}$ but not to $M^{(i)}$.

We shall see later⁴ that each of the bargaining sets $M_0^{(i)}$, $M^{(i)}$

¹If (2.1) is not assumed, we require $x_i \geq v(i)$ for all i .

²Example 3.1. In this example coalitional rationality coincides with individual rationality.

³A similar example was suggested to me by H. Kesten, in a different context.

⁴Two examples of this kind were indicated also in [1].

and $\mathcal{M}_1^{(i)}$ may be different from \mathcal{M}_0 , \mathcal{M} and \mathcal{M}_1 .

Theorem 5.3 Let Γ be an n-person game, in which the set of the permissible coalitions consists of the 1, n-1, and n-person coalitions. The bargaining sets $\mathcal{M}_0^{(i)}$, $\mathcal{M}^{(i)}$, and $\mathcal{M}_1^{(i)}$ are the same for Γ . Moreover, the p.c.'s in these bargaining sets which are not of the form $(x; N)$ coincide with those in \mathcal{M} .

Proof: Any i.r.p.c. not of the form $(x; N)$ must be also coalitionally rational. P.c.'s of the form $(x; N)$ never enter into objections or counter objections. This, and the fact that $\mathcal{M}_0 = \mathcal{M} = \mathcal{M}_1$, assures the last assertion.

Any objection in $(x; N)$, in any of the above definitions, must be of the form $(x; l, N^{(l)})$, and any counter objection to such an objection, in any of the above definitions, must either satisfy $x_l = 0$ or be of the form $(x; k, N^{(k)})$, where player k belongs to the objecting K . Thus, one sees easily that conditions that $(x; N)$ is $\mathcal{M}_0^{(i)}$, $\mathcal{M}^{(i)}$, $\mathcal{M}_1^{(i)}$ -stable are equivalent. This completes the proof.

We realize that the only new p.c.'s which might appear in $\mathcal{M}^{(i)}$ and not in \mathcal{M} are p.c.'s of the form $(x; N)$. We shall see that this will actually happen. It is remarkable, however, that $\mathcal{M}^{(i)}$ -stable p.c.'s of the form $(x; N)$ which are not \mathcal{M} -stable, occur if and only if N is not an effective coalition. In other words, as long as \mathcal{M} -stable p.c.'s of the form $(x; N)$ exist, all the bargaining sets considered here are identical for our games. If no \mathcal{M} -stable p.c. of this form exists, the players may relax their stability requirements, by replacing the coalitional rationality requirement by individual rationality restriction. Doing so, they will always find an $\mathcal{M}^{(i)}$ -stable p.c. of the form $(x; N)$.

In order to prove this and analyze the new p.c.'s, we shall first

establish the following:

Lemma 5.1 Let Γ be the game described in Theorem 5.3. A p.c. of the form $(x; N)$ is $\mathcal{M}^{(i)}$ stable, if and only if for each player ℓ , $\ell = 1, 2, \dots, n$,

$$(5.3) \quad \sum_{i \neq \ell} x_i \geq v^{(\ell)} \quad \text{or} \quad x_\ell = 0 \quad \text{or} \quad v^{(k)} + x_k \geq v^{(\ell)} + x_\ell \quad \text{for all } k .$$

Here, $v^{(v)} \equiv v(N^{(v)}) \equiv v(N - \{v\})$, $v = 1, 2, \dots, n$.

The proof of this lemma is similar to the proof of Lemma 4.6, and will therefore be omitted.

Theorem 5.4 Let Γ be an n -person game, $n \geq 2$, in which the set of permissible coalitions consists of the 1, $n-1$, and n -person coalitions. If N is an effective coalition, then a p.c. of the form $(x; N)$ is $\mathcal{M}^{(i)}$ -stable if and only if it is \mathcal{M} -stable.

Proof: If $(x; N)$ is \mathcal{M} -stable, then it is a c.r.p.c., and the first inequality in (5.3) holds for each ℓ . Hence $(x; N)$ is also $\mathcal{M}^{(i)}$ -stable.

Conversely, suppose that $(x; N)$ is $\mathcal{M}^{(i)}$ -stable. This means that (5.3) holds. If there exists a player μ for which

$$(5.4) \quad \sum_{i \neq \mu} x_i < v^{(\mu)} \quad \text{and} \quad x_\mu = 0 ,$$

then

$$(5.5) \quad v(N) = \sum_{i=1}^n x_i = \sum_{i \neq \mu} x_i < v^{(\mu)} .$$

By Corollary 4.1, N is not an effective coalition, contrary to our assumption. If there exists a player μ for which

$$(5.6) \quad \sum_{i \neq \mu} x_i < v^{(\mu)} \quad \text{and} \quad v^{(k)} + x_k \geq v^{(\mu)} + x_\mu \quad \text{for all } k ,$$

then

$$(5.7) \quad v(N) = \sum_{i=1}^n x_i < v^{(\mu)} + x_\mu \leq v^{(k)} + x_k \quad \text{for all } k .$$

Summing up these inequalities for $k = 1, 2, \dots, n$, we obtain:

$$(5.8) \quad (n-1)v(N) < \sum_{k=1}^n v^{(i)} .$$

Again, by Corollary 4.1, N is not an effective coalition, contrary to our assumption. Thus, the first inequality in (5.3) must hold for each l , whence, by Theorem 4.1, $(x; N)$ is \mathcal{M} -stable. This completes the proof.

The situation is more complicated if N is not an effective coalition. Several cases will be distinguished.

Theorem 5.5. Let Γ be an n -person game, $n \geq 3$, in which the set of the permissible coalitions consists of the 1, $n-1$, and n -person coalitions. Let N be a non-effective coalition. If the p.c.

$$(5.9) \quad (\omega_1 - c, \omega_2 - c, \dots, \omega_n - c; N) ,$$

where

$$(5.10) \quad c = \frac{\omega_1 + \omega_2 + \dots + \omega_n - v(N)}{n} ,$$

is individually rational¹, then this p.c. is $\mathcal{M}^{(i)}$ -stable, and no other p.c. of the form $(x; N)$ is $\mathcal{M}^{(i)}$ -stable. In particular, (5.9) is always individually rational if the characteristic function of the game Γ is super-additive.²

Proof: Case A. Suppose that

$$(5.11) \quad v(N) \geq v(N - \{v\}) \equiv v^{(v)} \quad \text{for each } v, v = 1, 2, \dots, n .$$

By Corollary 4.1, since N is not an effective coalition, we obtain:

$$(5.12) \quad (n-1)v(N) < \sum_{i=1}^n v^{(i)} .$$

Therefore,

$$(5.13) \quad (n-1)v^{(v)} < \sum_{i=1}^n v^{(i)} \quad \text{for all } v, v = 1, 2, \dots, n ,$$

¹I.e., if $\omega_v - c \geq 0$ for each $v, v = 1, 2, \dots, n$.

²I.e., if (5.11) holds.

which, by (4.8), implies

$$(5.14) \quad (n-1) \sum_{i \neq v} \omega_i < \sum_{i=1}^n \sum_{j \neq i} \omega_j = (n-1) \sum_{i=1}^n \omega_i .$$

We see that $\omega_v > 0$ for all v , $v = 1, 2, \dots, n$; hence,

$$(5.15) \quad \omega_v - c = \frac{n\omega_v + v(N) - \omega_1 - \omega_2 - \dots - \omega_n}{n} = \frac{(n-1)\omega_v + v(N) - v^{(v)}}{n} > 0 .$$

This proves that (5.9) is indeed an i.r.p.c. Since (5.9) is an i.r.p.c., then, by (5.3), it is $\mathcal{M}^{(i)}$ -stable, because for each k and ℓ ,

$$(5.16) \quad v^{(k)} + (\omega_k - c) = \sum_{i=1}^n \omega_i - c = v^{(\ell)} + (\omega_\ell - c) .$$

Conversely, let $(x; N)$ be an $\mathcal{M}^{(i)}$ -stable p.c. for Γ , then (5.3) is satisfied. Since, by Theorem 4.1, the game has an empty core, there exists a player $\ell = \mu$, for which the first inequality of (5.3) does not hold. We know that (5.4) implies (5.5), which is contrary to (5.11), hence we must conclude that $x_\mu > 0$, and that

$$(5.17) \quad v^{(k)} + x_k \geq v^{(\mu)} + x_\mu \text{ for all } k, \quad \sum_{i \neq \mu} x_i < v^{(\mu)} .$$

If, for some player ρ we had

$$(5.18) \quad \sum_{i \neq \rho} x_i \geq v^{(\rho)} ,$$

we would obtain

$$(5.19) \quad v(N) = \sum_{i=1}^n x_i \geq v^{(\rho)} + x_\rho \geq v^{(\mu)} + x_\mu > \sum_{i=1}^n x_i = v(N) ,$$

which is a manifest contradiction. Thus, (5.18) never holds, $x_v > 0$ for each player v ; and hence, by (5.3)

$$(5.20) \quad v^{(k)} + x_k = v^{(\ell)} + x_\ell, \quad \sum_{i=1}^n x_i = v(N)$$

holds for each k and ℓ . Equations (5.20) have a unique solution; therefore $(x; N)$ is the p.c. (5.9).

Case B. We now suppose that (5.11) does not hold. The fact that (5.9) is an i.r.p.c. is by (4.2) and (5.15) equivalent to

$$(5.21) \quad v(N) + \sum v^{(i)} \geq n v^{(v)} \quad \text{for all } v, v = 1, 2, \dots, n.$$

As before, (5.9) is certainly $\mathcal{M}^{(i)}$ -stable because (5.16) holds.

Let $(x; N)$ be an $\mathcal{M}^{(i)}$ -stable p.c. for Γ , then (5.3) is satisfied. Let S be the set of indices v , for which

$$(5.22) \quad \sum_{i \neq v} x_i < v^{(v)},$$

and let $R = N - S$. S is not empty, because the game has an empty core. Obviously,

$$(5.23) \quad v(N) = \sum_{i=1}^n x_i < v^{(v)} + x_v \quad \text{for each } v, v \in S,$$

$$(5.24) \quad v(N) = \sum_{i=1}^n x_i \geq v^{(\mu)} + x_\mu \quad \text{for each } v, v \in R.$$

I. We shall examine now the case in which $x_v = 0$ for each $v, v \in S$. In this case $v(N) < v^{(v)}$ for each $v, v \in S$. If¹ $|R| \neq 0$, we can sum up the inequalities (5.24), and we obtain

$$(5.25) \quad (|R| - 1) v(N) \geq \sum_{i \in R} v^{(i)}.$$

Let $v^{(\alpha)} = \text{Max}(v^{(1)}, v^{(2)}, \dots, v^{(n)})$. Certainly, $\alpha \in S$, because $v^{(\alpha)} > v(N)$. From (5.21) we now obtain

$$(5.26) \quad n v^{(\alpha)} \leq v(N) + \sum_{i \in S} v^{(i)} + (|R| - 1) v(N) \leq |R| v(N) + |S| v^{(\alpha)},$$

which implies $v^{(\alpha)} \leq v(N)$. This, however, cannot happen, because we are dealing with the case in which (5.11) does not hold.

Thus, $R = \emptyset$, hence $v(N) = 0$. In this case, $(x; N)$ must be $(0, 0, \dots, 0; N)$. We shall show that it must be equal to the p.c. (5.9). This follows easily from (5.10) and from the fact that (5.9) is an individually

¹ $|P|$ denotes the number of elements in a set P .

rational p.c.

II. Suppose that there exists a player ρ , $\rho \in S$, for which $x_\rho \neq 0$.

Certainly, $v(N) > 0$. By (5.3), then

$$(5.27) \quad v^{(k)} + x_k \geq v^{(\rho)} + x_\rho \quad \text{for each } k, k = 1, 2, \dots, n.$$

Again, $R = \emptyset$, since otherwise, for a player σ in R we would have obtained by (5.23), (5.24) and (5.27) the impossible inequality:

$$(5.28) \quad v(N) \geq v^{(\sigma)} + x_\sigma \geq v^{(\rho)} + x_\rho > v(N).$$

Thus, $S = N$. Let P be the set of players τ for which $x_\tau = 0$, and let $v^{(\alpha)} = \text{Max}(v^{(1)}, v^{(2)}, \dots, v^{(n)})$. Let Q be the set of players ρ for which $x_\rho > 0$. We know that $Q \neq \emptyset$ and that (5.27) holds for each ρ , $\rho \in Q$; hence

$$(5.29) \quad v^{(\tau)} \geq v^{(\rho)} + x_\rho \quad \text{whenever } \tau \in P, \rho \in Q.$$

Summing up these relations for all the ρ 's in Q , we obtain

$$(5.30) \quad |Q| v^{(\tau)} \geq \sum_{i \in Q} v^{(i)} + v(N) \quad \text{whenever } \tau \in P,$$

which implies

$$(5.31) \quad v(N) + \sum_{i \in Q} v^{(i)} + \sum_{i \in P} v^{(i)} \leq |Q| v^{(\tau)} + \sum_{i \in P} v^{(i)} \leq nv^{(\alpha)}.$$

By (5.21) we realize that this situation can happen only if $v^{(\tau)} = v^{(\alpha)}$ for each τ , $\tau \in P$, and if

$$(5.32) \quad v(N) + \sum v^{(i)} = nv^{(\tau)} \quad \text{whenever } x_\tau = 0.$$

It follows that $\omega_\tau - c = 0$ whenever $x_\tau = 0$. The inequalities (5.20), restricted to all k, ℓ , where $k, \ell \in Q$, determine uniquely x_ρ for each ρ , $\rho \in Q$, hence $(x; N)$ is the p.c. (5.9). This completes the proof of the theorem.

If (5.9) is not an i.r.p.c., it cannot be $\mathcal{M}^{(i)}$ -stable. In order to get the stable p.c.'s in this case, one has to refer to some sort of "deleting" of players from the game. This will be established as follows:

Lemma 5.2 Let Γ be a game as described in Theorem 5.5, except that (5.9) be no longer individually rational. Let α be ^{any} one of the players having the smallest quota, and let $(x; N)$ be an $\mathcal{M}^{(i)}$ -stable p.c. Under these conditions, $x_\alpha = 0$.

Proof: We know that in this case the characteristic function of the game is not superadditive (see end of Theorem 5.5). Certainly $\omega_\alpha - c < 0$.

If $x_\alpha > 0$, then, by (5.3),

$$(5.33) \quad \text{either } \sum_{i \neq \alpha} x_i \geq v^{(\alpha)} \quad \text{or} \quad v^{(k)} + x_k \geq v^{(\alpha)} + x_\alpha \quad \text{for each } k.$$

By (4.2), $v^{(\alpha)} = \text{Max} (v^{(1)}, v^{(2)}, \dots, v^{(n)})$; hence the first inequality in (5.33) implies $v(N) \geq v^{(\alpha)} + x_\alpha > v^{(\alpha)}$, which is impossible because (5.11) does not hold, and the second inequality in (5.33) implies

$$(5.34) \quad \sum_{i=1}^n v^{(i)} + v(N) \geq nv^{(\alpha)} + nx_\alpha > nv^{(\alpha)},$$

which is again impossible because it implies (see (5.21)) $\omega_\alpha - c > 0$.

Therefore $x_\alpha = 0$.

Theorem 5.6 Let Γ be an n-person game, $n > 3$, in which the set of permissible coalitions consists of the 1, n-1, and n-person coalitions. Let α be a player having the smallest quota, and let $\omega_\alpha - c < 0$, where c is defined by (5.10). Let Γ^* be an (n-1)-person game which results from Γ by deleting the player α and assigning the remaining players the following characteristic function:

$$(5.35) \quad v^*(B) = \begin{cases} v(B \cup \{\alpha\}) & \text{whenever } B = N - \{\alpha\} - \{v\}, v = 1, 2, \dots, n, v \neq \alpha, \\ v(N) & \text{for } B = N - \{\alpha\}, \\ 0 & \text{for } B = 1, 2, \dots, \alpha-1, \alpha+1, \dots, n. \end{cases}$$

Under these conditions,

$$(5.36) \quad (x; N) \equiv (x_1, x_2, \dots, x_n; N)$$

is $\mathcal{M}^{(i)}$ -stable with respect to Γ , if and only if $x_\alpha = 0$ and

$$(5.37) \quad (x^*; N^*) \equiv (x_1, x_2, \dots, x_{\alpha-1}, x_{\alpha+1}, \dots, x_n; N - \{\alpha\})$$

is $\mathcal{M}^{(i)}$ -stable with respect to Γ^* .

Proof: We know already that $x_\alpha = 0$ is a necessary condition for $(x; N)$ to be $\mathcal{M}^{(i)}$ -stable with respect to Γ , hence (5.3) reduces to

$$(5.38) \quad \sum_{i \neq v, \alpha} x_i \geq v^{(v)} \quad \text{or} \quad x_v = 0 \quad \text{or} \quad v^{(k)} + x_k \geq v^{(v)} + x_v \quad \text{for each } k,$$

which are to be satisfied for each $v, v \neq \alpha, v = 1, 2, \dots, n$. The corresponding conditions for $(x^*; N^*)$ to be $\mathcal{M}^{(i)}$ -stable with respect to Γ^* are the same inequalities (5.38), except that k is not required to cover player α . The theorem will be proved if we can show that no player φ exists, for which

$$(5.39) \quad v^{(k)} + x_k \geq v^{(\varphi)} + x_\varphi \quad \text{for each } k, k \neq \alpha, \quad v^{(\alpha)} + x_\alpha < v^{(\varphi)} + x_\varphi,$$

if $(x; N)$ is an i.r.p.c. with respect to Γ . Indeed, otherwise, we obtain

$$(5.40) \quad v^{(k)} + x_k > v^{(\alpha)} \quad \text{for each } k, k \neq \alpha,$$

and hence

$$(5.41) \quad \sum_{i=1}^n v^{(i)} + v(N) > nv^{(\alpha)}.$$

This contradicts the fact that $\omega_\alpha - c < 0$ (see (5.21)).

Discussion.

The $\mathcal{M}^{(i)}$ -stable p.c.'s of the form $(x; N)$ are the core of the game, provided that the core is not empty. If the core is empty but (5.9) is an i.r.p.c., it is also the unique $\mathcal{M}^{(i)}$ -stable p.c. of this form. If this is not the case, in order to obtain the $\mathcal{M}^{(i)}$ -stable p.c. of the form $(x; N)$, one "deletes" the "weakest" players one at a time, as explained in Theorem 5.6, until one arrives at a game for which the $\mathcal{M}^{(i)}$ -stable p.c.'s are known. The players which are deleted would get a zero payoff anyway, and the payoff for the others is determined to be their payoffs in the $\mathcal{M}^{(i)}$ -stable p.c.'s in the reduced game. Our analysis will be completed if we list the $\mathcal{M}^{(i)}$ -stable p.c.'s for the 3-person game with an empty core, for which (5.9) is not individually rational:

Theorem 5.7 Let Γ be a 3-person game, the characteristic function of which satisfies

$$(5.42) \quad v^{(1)} \geq v^{(2)} \geq v^{(3)}, \quad v(N) < v^{(1)}, \quad v(N) + v^{(2)} + v^{(3)} < 2v^{(1)},$$

where $v^{(i)} = v(N - \{i\}$, $i = 1, 2, 3$. The following are the necessary and sufficient conditions for a p.c. $(x; N) = (x_1, x_2, x_3; 123)$ to be $\mathcal{M}^{(i)}$ -stable with respect to Γ :

- (i) If $v(N) \geq v^{(2)} + v^{(3)}$ then $x_1 = 0$, $x_2 \geq v^{(3)}$, $x_3 \geq v^{(2)}$ and
 $x_2 + x_3 = v(N)$.
- (ii) If $v(N) < v^{(2)} + v^{(3)}$ but $v(N) + v^{(3)} \geq v^{(2)}$ and $v(N) + v^{(2)} \geq v^{(3)}$,
then $x_1 = 0$, $x_2 = \frac{1}{2}(v(N) + v^{(3)} - v^{(2)})$, $x_3 = \frac{1}{2}(v(N) + v^{(2)} - v^{(3)})$.
- (iii) If $v(N) + v^{(3)} - v^{(2)} < 0$, then $x_1 = x_2 = 0$, $x_3 = v(N)$.

The proof follows immediately from analyzing the inequalities (5.3).

Corollary 5.1. For our games, there always exists an $\mathcal{M}^{(i)}$ -stable p.c. of the form $(x; N)$.

Corollary 5.2 The procedure described in Section 4, of deleting a weak player from the game in order to find the \mathcal{M} -stable p.c.'s of the form $(x; j, N^{(j)})$, is the same procedure for finding the $\mathcal{M}^{(i)}$ -stable p.c.'s of the same form. In other words, when we treat the reduced game, we may look for the $\mathcal{M}^{(i)}$ -stable p.c.'s for this game, and forget about the \mathcal{M} -stability theory. This is a consequence of the fact that the reduced game always has a non-empty core (see end of case (i) in Theorem 4.3 and the conclusion at the end of Section 4). For games with a non-empty core, $\mathcal{M} = \mathcal{M}^{(i)}$.

Princeton University and

The Hebrew University of Jerusalem

REFERENCES

- [1] R. J. Aumann and M. Maschler. The bargaining set for cooperative games. (Expected to appear in No. 52 of Annals of Mathematics Studies, Princeton University Press, Princeton, New Jersey.)
- [2] G. K. Kalisch. Generalized quota solutions of n-person games. Contributions to the Theory of Games, Vol. IV, Annals of Mathematics Studies, No. 40, Princeton University Press, Princeton, New Jersey (1959), pp. 163-177.
- [3] M. Maschler. Stable payoff configurations for quota games. (To appear.)
- [4] L. S. Shapley. Quota solutions of n-person games. Contributions to the Theory of Games, Vol. II, Annals of Mathematics Studies, No. 28, Princeton University Press, Princeton, New Jersey (1953), pp. 343-359.